Disease mechanisms and neuroprotection by tauroursodeoxycholic acid in Rpgr knockout mice

Zhang, Xun; Shahani, Uma; Reilly, James; Shu, Xinhua

Published in:
Journal of Cellular Physiology

DOI:
[10.1002/jcp.28519](https://doi.org/10.1002/jcp.28519)

Publication date:
2019

Document Version
Author accepted manuscript

[Link to publication in ResearchOnline](#)

Citation for published version (Harvard):
Zhang, X, Shahani, U, Reilly, J & Shu, X 2019, 'Disease mechanisms and neuroprotection by tauroursodeoxycholic acid in Rpgr knockout mice', *Journal of Cellular Physiology*, vol. 234, no. 10, pp. 18801-18812. <https://doi.org/10.1002/jcp.28519>

General rights

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

Take down policy

If you believe that this document breaches copyright please view our takedown policy at <https://edshare.gcu.ac.uk/id/eprint/5179> for details of how to contact us.

Disease mechanisms and neuroprotection by tauroursodeoxycholic acid in *Rpgr* knockout mice

Xun Zhang¹, Uma Shahani², James Reilly¹, Xinhua Shu^{1,2*}

1 Department of Biological and Biomedical Sciences, Glasgow Caledonian University, Cowcaddens Road, Glasgow G4 0BA

2 Department of Vision Science, Glasgow Caledonian University, Cowcaddens Road, Glasgow G4 0BA

* Corresponding author Dr Xinhua Shu, Email Xinhua.Shu@gcu.ac.uk

Abstract

Mutations in the *RPGR* gene are the predominant cause of retinitis pigmentosa (RP). RPGR plays a critical role as a scaffold protein in the regulation of protein trafficking from the basal body to the axoneme, where the cargoes are transported to the outer segments (OS) of photoreceptors. This trafficking process is controlled directly by intraflagellar transport (IFT) complexes and regulated by the RPGR protein complex, although the precise mechanisms have yet to be defined. We employed an *Rpgr* conditional knockout (cko) mouse model to investigate the disease mechanisms during retinal degeneration and to evaluate the protective effects of tauroursodeoxycholic acid (TUDCA). Rhodopsin, cone opsins and transducin were mislocalized in *Rpgr* cko photoreceptors, while localization of NPHP4 to connecting cilia was absent, suggesting that RPGR is required for ciliary protein trafficking. Microglia were activated in advance of retinal degeneration in *Rpgr* cko mouse retinas. TUDCA treatment suppressed microglial activation and inflammation and prevented photoreceptor degeneration in *Rpgr* cko mice. Our data demonstrated that TUDCA has therapeutic potential for RPGR-associated RP patients.

KEYWORDS retinitis pigmentosa, RPGR, microglia activation, tauroursodeoxycholic acid, neuroprotection

1 INTRODUCTION

Retinitis pigmentosa (RP) is a class of inherited retinal disorder that causes progressive visual impairment and which can lead ultimately to blindness. The worldwide prevalence of RP is about 1 in 4,000, meaning there are more than 1 million people currently living with RP (Hartong et al., 2006). The disorder is characterised by pigmentation in retinal cells that is caused by deposition of materials from retinal pigment epithelium (RPE) cells and degeneration of photoreceptors. In most RP cases, patients lose their peripheral vision and night vision, a process often beginning in adolescence and caused by the death of rod photoreceptors. The subsequent degeneration of cone photoreceptor in later stages results in loss of central vision and colour perception. Some patients, on the other hand, exhibit cone-rod dystrophy, which involves an initial loss of cone photoreceptors followed by degeneration of rods (Hamel, 2007).

Mutations in the retinitis pigmentosa GTPase regulator (*RPGR*) genes are the major single cause of RP, accounting for up to 20% of cases in Caucasians (Shu et al., 2007). The *RPGR* gene has more than 10 alternative transcripts, of which *RPGR^{ex1-19}* and *RPGR^{ORF15}* are well-studied major transcripts (Shu et al., 2005; Vervoort et al., 2000). The *RPGR^{ex1-19}* transcript has been shown to be widely expressed in all examined tissues in different species, including human, mouse, *Xenopus* and zebrafish, whereas the *RPGR^{ORF15}* transcript is predominantly expressed in the retina and contains a mutation hot spot in the C-terminal exon, called ORF15 (Shu et al., 2006; Raghupathy et al., 2015). The *RPGR^{ex1-19}* isoform is conserved in both vertebrates and invertebrates, whereas, the *RPGR^{ORF15}* isoform is unique to vertebrates (Raghupathy et al., 2015). *RPGR* forms a complex with other proteins such as *RPGR*-interacting protein 1 (*RPGRIP1*) and *RPGRIP1*-like protein (*RPGRIP1L*) in the connecting cilium to regulate transport of cargoes such as rhodopsin (Patnaik et al., 2015). Knockout (ko) of *Rpgr* in mice caused a reduced level of rhodopsin, partially mislocalized blue and green cone opsins, and notable abnormality of newly formed disk membranes at the base of photoreceptors (Hong et al., 2000). Interestingly, significantly increased rhodopsin was mistrafficked in the ventral retina compared to the dorsal retina; however, there was more mistrafficked green opsin in the dorsal retina than in the ventral retina (Charng et al., 2016). Partial mislocalization of rhodopsin and opsin has also been reported in the retinas of *XLPR2* dogs, which carry a two-nucleotide deletion in the *RPGR*

exon ORF15 and which exhibited early-onset retinal degeneration (Beltran et al., 2006). *Rpgr* ko mice were found to exhibit moderate degeneration with approximately 25% loss of outer nuclear layer (ONL) at 6 months of age (Hong et al., 2000). *Rpgr* conditional knockout (cko) mice also showed progressive photoreceptor degeneration with 65% loss of ONL at ~13 months of age (Huang et al., 2012).

In this study, we investigated photoreceptor death mechanisms and evaluated the neuroprotective effects of tauroursodeoxycholic acid (TUDCA) in *Rpgr* cko mice. We found that cko mouse retinas had significant increases in cell death and inflammation when compared to wildtype mouse retinas. TUDCA treatment resulted in decreased photoreceptor death and inhibited inflammation in *Rpgr* cko mouse retinas.

2 MATERIALS AND METHODS

2.1 Animals and TUDCA administration

Rpgr cko mouse (Huang et al., 2012) were gifted from Professor Wright's lab at MRC Human Genetics Unit, Edinburgh, and bred in the Animal Unit at Glasgow Caledonian University under a 14:10 hour light-dark cycle. Genotyping of *Rpgr* cko mice was performed by PCR and sequencing. Both wildtype siblings and *Rpgr* cko mice (male and female) were used for this study. Previous studies have shown that injection of TUDCA at dose 500mg/kg is effective in preventing retinal degeneration in RP mouse models (Drack et al., 2012); consequently, this dose was chosen for our study. Six *Rpgr* cko mice were injected weekly with TUDCA in 0.15M NaHCO₃ at 500mg/kg dose intraperitoneally from postnatal days (P) 30 to P120 (n=6). Untreated control mice received the same volume of 0.15M NaHCO₃ (n=6). The body weight of mice in each group was measured prior to injection each week. There was no significant difference in body weight growth between untreated and TUDCA-treated group (data not shown). The animal experiment was approved by the Animal Ethics and Welfare Committee, Department of Life Sciences, Glasgow Caledonian University in accordance with the UK home office animal care guidelines (Project licence P8C815DC9).

2.2 Histology and immunohistochemistry

Enucleated eyes were marked with marker pen to distinguish superior and inferior side and were fixed in 2% PFA/PBS at 4°C for 18-24 hours. The fixed whole eyes then went through 10%, 30%,

50%, 70%, 90%, 100%×3 ethanol each for 1h, followed by two changes of Histo-Clear (Sigma, USA) and three changes of paraffin in 60°C oven. Next, eyes were embedded in paraffin and completely sectioned at 8µm thickness through the vertical meridian. The paraffin sections (within the optic nerve head region) were gradually rehydrated by undergoing two changes of Histo-Clear, two changes of 100% ethanol (5min/each), 90% ethanol for 2min, 70% ethanol for 2min, 50% ethanol for 2min and distilled water for 2min before staining with hematoxylin (Sigma, USA) for 8 min. Slides were then washed in running tap water for 20 min and dehydrated by going through 50% ethanol for 2min and 70% ethanol for 2 min, then counterstained with eosin (Sigma, USA) for 1min, followed by further dehydration with 90% ethanol for 1min, two changes of 100% ethanol (5min/each) and two changes of Histo-Clear (5min/each). Slides were examined and photographed under a light microscope (Olympus, Japan). For measurement of the outer nuclear layer (ONL), two retinal sections of each eye were selected and images were taken from superior and inferior sides of the optic nerve head (ONH); ONL thickness was measured at intervals of 0.2mm. Five eyes from five individual mice from each group were used for ONL measurement.

Fixed eye samples were processed through a series of sucrose concentrations (5%, 15% and 20%) for 4 hours each step, followed by embedding in OCT (optimal cutting temperature) compound and subsequent freezing. Eyes were completely sectioned at 10µm thickness through the vertical meridian. The sections were rehydrated with wash buffer (1×TBS/0.025% Triton X-100) twice (5min/each). The sections were then incubated with blocking buffer (1×TBS/0.3% Triton X-100/5% sheep serum) for 1 hour at room temperature and incubated with primary antibodies (Table S1) in blocking buffer overnight at 4°C. Sections were washed 3 times (5min/each) and incubated with FITC-conjugated AF488 or AF594 secondary antibody (1:400 dilution) (Thermo Fisher Scientific, UK) for 1 hour at room temperature. Tissue sections were counterstained with DAPI (Thermo Fisher Scientific, UK) after washing 5 times (5min/each) with wash buffer. Fluorescence images were captured on a ZEISS LSM800 microscope (Zeiss, Germany). To measure the intensity of the fluorescence signal, two retinal sections of each eye were collected: one region (10µm×10µm, under 400x magnification) in the superior side (0.4mm from the ONH) and one region (10µm×10µm, under 400x magnification) in the inferior side (0.4mm from

ONH) in each section were chosen for fluorescent signal quantification. The intensities of fluorescent signals from the superior region and from the inferior region were averaged for final calculation. Five eyes from each group were used for the quantification.

2.3 Quantitative real-time polymerase chain reaction (qRT-PCR)

qRT-PCR was performed with PCR-Platinum® SYBR® Green qPCR SuperMix- UDG kit (Invitrogen). Briefly, the reactions were set up in a 96-well plate as follows: each 15µl reaction contained 7.5µl platinum SYBR green qPCR SuperMix-UDG with ROX, 0.4µl 10µM of forward and reverse primers, 3µl diluted cDNA sample (50ng/µl) and 5.7µl of nuclease-free water. The no template controls (NTC), which contained all the components except for the cDNA template, were also set up and topped up to 15µl with nuclease-free water. DNA amplification and fluorescence detection were performed according to previous description (Tohari et al., 2016). Sequences of primers used for qRT-PCR are *Caspase 3* forward 5' TGGTGATGAAGGGGTCATTTATG 3' and reverse 5' TTCGGCTTTCCAGTCAGACTC 3'; *IL-β* forward 5' GGAGAACCAAGCAACGACAAAATA 3' and reverse 5' TGGGGAAGTCTGCAGACTCAAAC 3'; *Gapdh* forward 5' GTCTCCTGCGACTTCAGC 3' and reverse 5' TCATTGTCATACCAGGAAATGAGC 3'.

2.4 Western blotting

The retinas were lysed and proteins were extracted using T-PER™ Tissue Protein Extraction Reagent (Thermo Fihser Scientific, UK), and the concentration of protein lysis was measured by protein assay (Biorad, USA). Lysis were incubated with 4×loading buffer (Thermo Fisher Scientific, UK) at 70°C for 10 min before loading. Proteins were separated with precast gel (BioRad, UK) and transferred to the nitrocellulose membrane (GE Healthcare, UK). The membrane was blocked in 5% milk for 1h and incubated with primary antibodies at 4°C for about 18 hours. The IRDye secondary antibody (Li-cor, USA) was used for detection of targeting protein.

2.5 TUNEL assay

The mouse eyes were fixed and cut as described above. The cryosections (8µm) of mouse eyes were used to perform a TUNEL (Terminal deoxynucleotidyl transferase dUTP nick end labelling) assay to detect apoptotic cells by the DeadEnd™ Fluorometric TUNEL System (Promega, UK)

according to the manufacturer's instruction. Samples were washed with 1×PBS for 5 min before being fixed by 4% PFA/PBS and permeabilized with 20µg/ml Proteinase K solution. Samples were then treated with rTdT Incubation Buffer for 1 hour at 37°C and the reaction was stopped by 2×SSC. Slides were mounted with DAPI stain and the fluorescence was photographed by ZEISS LSM 800.

2.6 Statistical analysis

All data are presented as mean ± SD. Data were analysed and compared between two groups at different time-points by a non-parametric *t*-test following by Wilcoxon matched-pairs signed rank test with GraphPad Prism software. Data from TUDCA treatment were analysed using one-way ANOVA followed by Bonferroni test. Differences were regarded as statistically significant if $p < 0.05$ and are denoted by asterisks * (* $p < 0.05$; ** $p < 0.01$; *** $p < 0.001$; **** $p < 0.0001$).

3 RESULTS

3.1 Abnormal ciliary trafficking in *Rpgr* cko mouse retinas

Using Sanger sequencing we found that a total of 3189 bp was deleted in the *Rpgr* cko mice, including proximal promoter, exons 1-3, and resulted in absence of RPGR expression and localization to connecting cilia of cko mouse photoreceptors (data not shown). To investigate the ciliary trafficking of phototransduction components in the *Rpgr* cko retina, immunohistochemistry of eye cryosections from one month-old mice was performed with anti-opsins, anti-transducin and anti-GRK1 antibodies. Rhodopsin was localised to outer segments of WT photoreceptors; in *Rpgr* cko mouse photoreceptor the rhodopsin was partially mislocalized to the connecting cilium. Similarly, red/green opsin and blue opsin were mistrafficked and accumulated in the connecting cilium, the perinuclear space and the outer plexiform layer (Figure 1). However, there was no difference in GRK1 localization in WT and *Rpgr* cko photoreceptors (data not shown).

NPHP4 is a component of RPGR protein complex by direct interaction with RPGRIP1 (Patnaik et al., 2015; Roepman et al., 2005). Mutations in *NPHP4* caused nephronophthisis and Senior-Løken syndrome (a combination of nephronophthisis and RP) (Mollet et al., 2012; Otto et al., 2002). NPHP4 is localized to the connecting cilium of photoreceptors, a localization that is dependent on RPGRIP1 (Patil et al., 2012; Roepman et al., 2005; Won et al., 2011). We performed immunohistochemistry in

paraffin sections of retina to detect NPHP4 localization. In WT retina, NPHP4 was present in the connecting cilium along with acetylated α tubulin. However, in *Rpgr* cko retina NPHP4 was absent from the connecting cilium (Figure 2A). We also examined NPHP4 expression in retinas by Western blotting and found that NPHP4 expression was significantly decreased at aged 3 months and above (Figure 2B,C; Figure S1).

3.2 Photoreceptor cell degeneration in *Rpgr* cko mice

To assess photoreceptor cell degeneration throughout ageing, we performed hematoxylin & eosin staining with paraffin sections of WT and *Rpgr* cko mouse eyes at ages 1, 3, 6, and 12 months and measured the thickness of the outer nuclear layer (ONL) at five different points on both superior and inferior sides of the retina. ONL thickness started to show a significant decrease in 3-month old *Rpgr* cko mice and was further markedly decreased at 6 months old, compared to age-matched WT mice. Only 2-3 layers of photoreceptor nuclei remained in 12-month old *Rpgr* cko mice (Figure S2). We also measured the length of photoreceptor outer segments in WT and *Rpgr* cko mice at different age points. At one-month old, the length of OSs was similar between WT and *Rpgr* cko mice; from the age of 3 months, however, OSs in cko mice were significantly shorter than those of WT mice (Figure S3).

Decreased thickness of the ONL in *Rpgr* cko mice was surmised to be due to death of photoreceptor cells. We performed a TUNEL assay to detect the total photoreceptor death at different age points. Significantly increased photoreceptor cell death was present in cko mouse retinas from 3 months of age and peaked at 6 months. In the age-matched WT retina, no significant cell death was observed (Figure 3). To determine whether photoreceptor cell death is caspase-dependent, we employed immunohistochemistry to detect activation of Caspase-3, the key component of the caspase-dependent cell death pathway, and found that cleaved caspase-3 was present in the outer plexiform layer (OPL) and ONL at age 3, 6 and 12 months (Figure S4).

3.3 Microglia were activated at the early stage of retinal degeneration

Microglia are immune cells resident in the central nervous system, including the retina (Li et al., 2015). In the healthy mammalian retina, microglia are in resting form, releasing anti-inflammatory factors to maintain homeostasis of the retina; however, they can be activated to the ramified form to

mediate phagocytosis in disease conditions (Karlstetter et al., 2015). To investigate if there was microglial activation in *Rpgr* cko mouse retinas, Iba-1, a biomarker of microglia, was measured by immunohistochemistry in cryosections from WT and *Rpgr* cko mouse eyes. In the latter the microglia were activated and migrated into the ONL as early as 1 month old, whereas in the age-matched WT retina they were present in the OPL (Figure 4).

To explore the consequence of microglial activation in *Rpgr* cko mouse retina, we examined the microglia-mediated inflammation. In macrophages or dendritic cells, the inflammasomes are assembled by scaffold protein NLRP3 with ASC adaptor and caspase-1 via PYD and CARD domain (Schroder & Tschopp, 2010). In the *Rpgr* cko retina, we found NLRP3 colocalized with Iba-1 in the ONL (Figure 5), indicating the formation of inflammasomes in activated microglia. The result suggests that microglia-mediated inflammation might contribute to the retinal degeneration in *Rpgr* cko mice.

3.4 TUDCA treatment ameliorated retinal degeneration in *Rpgr* cko mice

Tauroursodeoxycholic acid (TUDCA), the bile acid, has been widely used as an anti-apoptotic, anti-inflammatory and antioxidant compound (Pardue & Allen, 2018). TUDCA has been demonstrated to cause inhibition of retinal microglia activation and preservation of retinal structure and visual function (Pardue & Allen, 2018). To test the protective efficacy of TUDCA in *Rpgr* cko retina, *Rpgr* cko mice were intraperitoneally injected weekly with TUDCA (500mg/kg in 0.15M NaHCO₃) from P30 to P120 (n=6). Untreated control *Rpgr* cko mice received an equivalent volume of 0.15M NaHCO₃ (n=6). The thickness of the ONL was significantly increased in TUDCA-treated *Rpgr* cko mice when compared to untreated *Rpgr* cko mice (Figure 6A). The number of photoreceptors was significantly decreased in cko mice when compared to the wildtype mice; TUDCA treatment resulted in a higher number of photoreceptors compared to untreated cko mice (Figure S5). The decreased number of photoreceptor cells is possibly due to cell death; to investigate this we carried out a TUNEL assay and found that the number of photoreceptors undergoing cell death was notably reduced in the treated retina (Figure 6B). The caspase-dependent apoptosis pathway in cko mouse retina was possibly inhibited by TUDCA treatment, since expression of caspase3 on both protein and mRNA levels was notably decreased (Figure 6C, Figure S6A). In addition, TUDCA treatment significantly ameliorated

microglia activation by reducing the infiltration of activated microglia into the ONL (Figure 7A). The inflammasome formation in activated microglia was also suppressed (Figure 7B), followed by decreased expression of matured inflammatory cytokine IL-1 β (Figure 7C, Figure S6B).

4 DISCUSSION

In the current study we investigated mistrafficking of photoreceptor proteins, photoreceptor cell death and microglial activation, and evaluated the protective effects of TUDCA in the retina of *Rpgr* cko mice. We found that rhodopsin was partially mislocalized to the connecting cilia and that cone opsins were also partially mistrafficked to the cone inner segment, cell body and synapse (Figure 1). Partial mislocalization of rhodopsin and cone opsins in photoreceptors has also been reported in *Rpgr* ko mice (Hong et al., 2000; Charng et al., 2016). α -transducin, which operates in rod visual function, was also partially mislocalized in *Rpgr* cko mice. In fact, mislocalization of rhodopsin, cone opsins and α -transducin has also been reported in *Rrgrip1* ko mice and mutant zebrafish (Raghupathy et al., 2017; Won et al., 2009). NPHP4 is mutated in nephronophthisis with vision defects (Mollet et al., 2002) and directly interacts with RPGRIP1. Loss of NPHP4 or RPGRIP1 in mice resulted in abnormal development of outer segments (Won et al., 2009; 2011). Ptil et al. (2012) reported loss of NPHP4 localization to connecting cilia in RPGRIP1 ko mice. We also found that NPHP4 lost localization to connecting cilia in *Rpgr* cko mice and was decreased in cko retinas from 3 months old (Figure 2), possibly due to loss of photoreceptors, since photoreceptor death in cko mice was noticed from 3 months old and above. A proteomic study by Rao et al (2015) reported that proteins involved in ubiquitin-proteasome system or cilia function were decreased in *Rpgr* ko photoreceptor cilia and that some of these proteins were partially mislocalized (Rao et al., 2015). Our recent study also showed that RPGR protein complex regulated proteasome activities. These data suggest that the RPGR protein complex functions in ciliary protein trafficking and in maintenance of photoreceptor structure and function.

Microglia have been reported to mediate retinal degeneration (Karlstetter et al., 2015). Activated microglia have been detected in the retinas of RP patients (Gupta et al., 2003). Microglial activation has also been observed in preclinical RP rodent models including Pde6-alpha (*Pde6a*) and Pde6-beta (*Pde6b*) mutant mice, *Cngb1* knockout mice, homozygous P23H rats and Royal College of Surgeons

rats (Blank et al., 2018; Peng et al., 2014; Roche et al., 2016; Roque et al., 1996; Yoshida et al., 2013; Zeiss and Johnson, 2004; Zeng et al., 2005; Zhang et al., 2018; Zhao et al., 2015). In fact, microglia activation occurs prior to the initiation of photoreceptor degeneration. In rd10 mice, retinal microglial activation was initiated at postnatal day (P) 16 while the photoreceptor apoptosis started at P19 (Peng et al., 2014); a recent report showed that microglia were activated as early as P5 in rd10 mouse retinas (Roche et al., 2016). Early microglial activation ahead of photoreceptor cell death has also been reported in rd1 and *Cngb1* knockout mice (Blank et al., 2018; Zeiss et al., 2004). We observed early microglial activation in *Rpgr* cko mouse retinas at the age of one month when photoreceptor death had yet to be initiated (Figures 3 and 4). The data demonstrate that early microglial activation in retinas is a general feature of inherited retinal degeneration, possibly independent of genetic causes. The initiation of microglial activation in inherited retinal degeneration is not clear; it is possibly induced by toxic factors from pre-apoptotic photoreceptors. When photoreceptor death was halted by a genetic rescue, the activated microglia disappeared in photoreceptor degeneration sites and possibly became ramified (Zhang et al., 2018), indicating that photoreceptor death regulates microglial activation. It is presumed that activated microglia phagocytose pre-apoptotic mutant photoreceptors and secrete proinflammatory cytokines such as TNF- α and IL-1 β , which accelerate retinal degeneration (Karlstetter et al., 2015). Zhao et al. (2015) reported that both genetic depletion of microglia and suppression of microglial phagocytosis slowed rod cell death in rd10 mice. They also observed that photoreceptor cell death was suppressed following inhibition of IL-1 β signalling using an IL-1 receptor antagonist; IL-1 β was also significantly lower in mouse retina with genetic depletion of microglia when compared to control mice (Zhao et al., 2015). An early study reported that TNF- α expression was predominantly upregulated in activated microglia in rd mice (Zeng et al., 2005). We also found inflammasome formation and predominant expression of IL-1 β in activated microglia in *Rpgr* cko mouse retina (Figures 6 and 7). These data suggest that activated microglia-mediated-inflammation is involved in photoreceptor degeneration.

TUDCA has shown protective effects in a wide range of diseases including liver disease, kidney stones and gallstones, cardiovascular disease, diabetes, and neurodegenerative diseases (Pardue and Allen, 2018). TUDCA has also been used to treat rodent models of inherited retinal degeneration

including rd1, rd10, rd16, Bardet-Biedl syndrome 1 and *Lrat*^{-/-} mice, and transgenic Rhodopsin P23H rats; treated animals showed well-preserved retinal structure and improved visual function (Boatright et al., 2006; Drack et al., 2012; Fernandez-Sanchez et al., 2011; Noailles et al., 2014; Phillips et al., 2008; Zhang et al., 2012). It is proposed that protection of photoreceptor death by TUDCA treatment is due to improved protein folding and trafficking, reduced oxidative and ER stress, suppression of inflammation, decreased apoptosis, and increased RPE phagocytosis (Pardue and Allen, 2018). A recent report demonstrated that TUDCA inhibited microglial activation and decreased microglial distribution in outer retinal layers in *Rhodopsin* P23H homozygous rats (Noailles et al., 2014). We found that TUDCA treatment preserved the retinal structure of *Rpgr* cko mice (Figure 6), possibly through inhibition of apoptotic cell death since the number of apoptotic cells was significantly decreased in TUDCA-treated *Rpgr* cko mice when compared to untreated controls and since expression of the key component of apoptotic pathway, Caspase 3, was markedly decreased in TUDCA-treated *Rpgr* cko mouse retina (Figure 7). Zhang et al. (2012) also showed that TUDCA treated *Lrat*^{-/-} mice had decreased apoptosis with disappearance of activated Caspase 3. We also found that TUDCA treatment inhibited microglial activation and suppressed microglial migration to outer retinal layers (Figure 7). TUDCA treatment also reduced the NLRP3 inflammasome formation in activated microglia and decreased IL-1 β expression in treated *Rpgr* cko mouse retinas (Figure 7), which is consistent with previous studies that showed that TUDCA inhibited expression of inflammatory factors and promoted anti-inflammatory transcripts in rat microglia (Yanguas-Casás et al., 2017).

In conclusion, we observed defective ciliary protein trafficking and early microglial activation in the retinas of *Rpgr* cko mice. TUDCA treatment inhibited microglial activation and inflammation, resulting in preservation of retinal structure in *Rpgr* cko mice.

ACKNOWLEDGEMENTS

XZ was funded by a Fight for Sight PhD studentship (1419/20). We would like to thank the Rosetrees Trust (M160, M160-F1, M160-F2) and National Eye Research Centre (SCIAD063, SAC037) for supporting this work.

CONFLICTS OF INTEREST

The authors declare there are no conflicts of interest.

AUTHOR CONTRIBUTIONS

XZ performed the experiments. XS and US supervised the project. XZ, JR and XS analysed the data and wrote the manuscript.

REFERENCES

- Appelbaum, T., Becker, D., Santana, E., & Aguirre, G. D. (2016). Molecular studies of phenotype variation in canine RPGR-XLPRA1. *Molecular Vision*, 22, 319-331.
- Beltran, W.A., Hammond, P., Acland, G. M., & Aguirre, G. D. (2006) A frameshift mutation in RPGR exon ORF15 causes photoreceptor degeneration and inner retina remodeling in a model of X-linked retinitis pigmentosa. *Investigative Ophthalmology & Visual Science*, 47, 1669-1681.
- Blank, T., Goldmann, T., Koch, M., Amann, L., Schön, C., Bonin, M., ... Michalakis, S. (2018). Early microglia activation precedes photoreceptor degeneration in a mouse model of CNGB1-linked retinitis pigmentosa. *Frontiers in Immunology*, 8, 1930.
- Boatright, J. H., Moring, A. G., McElroy, C., Phillips, M. J., Do, V.T., Chang, B., ... Pardue, M. T. (2006). Tool from ancient pharmacopoeia prevents vision loss. *Molecular Vision*, 12, 1706–1714.
- Charng, J., Cideciyan, A. V., Jacobson, S. G., Sumaroka, A., Schwartz, S. B., Swider, M., ... Swaroop, A. (2016) Variegated yet non-random rod and cone photoreceptor disease patterns in RPGR-ORF15-associated retinal degeneration. *Human Molecular Genetics*, 25, 5444-5459.
- Drack, A.V., Dumitrescu, A.V., Bhattarai, S., Gratie, D., Stone, E. M., Mullins, R., & Sheffield, V.C. (2012). TUDCA slows retinal degeneration in two different mouse models of retinitis pigmentosa and prevents obesity in Bardet-Biedl syndrome type 1 mice. *Investigative Ophthalmology & Visual Science*, 53, 100-106.
- Fernandez-Sanchez, L., Lax, P., Pinilla, I., Martin-Nieto, J., & Cuenca, N. (2011). Tauroursodeoxycholic acid prevents retinal degeneration in transgenic P23H rats. *Investigative Ophthalmology & Visual Science*, 52, 4998–5008.

364 Gupta, N., Brown, K. E., & Milam, A. H. (2003). Activated microglia in human retinitis pigmentosa,
365 late-onset retinal degeneration, and age-related macular degeneration. *Experimental Eye Research*, 76,
366 463-471.

367 Hamel, C. P. (2007) Cone rod dystrophies. *Orphanet Journal of Rare Disease*, 2, 7.

368 Hartong, D. T., Berson, E. L., & Dryja, T. P. (2006) Retinitis pigmentosa. *Lancet*, 368, 1795-809.

369 Hong, D. H., Pawlyk, B. S., Shang, J., Sandberg, M. A., Berson, E. L., & Li, T. (2000) A retinitis
370 pigmentosa GTPase regulator (RPGR)-deficient mouse model for X-linked retinitis pigmentosa
371 (RP3). *Proceedings of the National Academy of Sciences of the United States of America*, 97, 3649–
372 3654.

373 Huang, W. C., Wright, A. F., Roman, A. J., Cideciyan, A. V., Manson, F. D., Gewaily, D. Y., ...
374 Jacobson, S. G. (2012) RPGR-associated retinal degeneration in human X-linked RP and a murine
375 model. *Investigative Ophthalmology & Visual Science*, 53, 5594–5608.

376 Karlstetter, M., Scholz, R., Rutar, M., Wong, W. T., Provis, J. M., & Langmann, T. (2015) Retinal
377 microglia: just bystander or target for therapy? *Progress in Retinal and Eye Research*, 45, 30-57.

378 Li, L., Eter, N., & Heiduschka, P. (2015) The microglia in healthy and diseased retina. *Experimental*
379 *Eye Research*, 136, 116–130.

380 Mollet, G., Salomon, R., Gribouval, O., Silbermann, F., Bacq, D., Landthaler, G., ... Saunier, S.
381 (2002) The gene mutated in juvenile nephronophthisis type 4 encodes a novel protein that interacts
382 with nephrocystin. *Nature Genetics*, 32, 300-305.

383 Noailles, A., Fernandez-Sanchez, L., Lax, P., & Cuenca, N. (2014). Microglia activation in a model of
384 retinal degeneration and TUDCA neuroprotective effects. *Journal of Neuroinflammation*, 11, 186.

385 Otto, E., Hoefele, J., Ruf, R., Mueller, A. M., Hiller, K. S., Wolf, M. T., ... Hildebrandt, F. (2002) A
386 gene mutated in nephronophthisis and retinitis pigmentosa encodes a novel protein, nephroretinin,
387 conserved in evolution. *American Journal of Human Genetics*, 71, 1161-1167.

388 Pardue, M. T., & Allen, R. S. (2018) Neuroprotective strategies for retinal diseases. *Progress in*
389 *Retinal and Eye Research*, 65, 50-76.

390 Patil, H., Tserentsoodol, N., Saha, A., Hao, Y., Webb, M., & Ferreira, P. A. (2012) Selective loss of
391 RPGRIP1-dependent ciliary targeting of NPHP4, RPGR and SDCCAG8 underlies the degeneration of
392 photoreceptor neurons. *Cell Death & Disease*, 3, e355.

393 Patnaik, S. R., Raghupathy, R. K., Zhang, X., Mansfield, D., & Shu, X. (2015) The Role of RPGR
394 and Its Interacting Proteins in Ciliopathies. *Journal of Ophthalmology*, 2015, 414781.

395 Patnaik, S. R., Zhang, X., Biswas, L., Akhtar, S., Zhou, X., Kusuluri, D. K., ... Shu, X. (2018) RPGR
396 protein complex regulates proteasome activity and mediates store-operated calcium entry. *Oncotarget*,
397 9, 23183-23197.

398 Peng, B., Xiao, J., Wang, K., So, K. F., Tipoe, G. L., & Lin, B. (2014) Suppression of microglial
399 activation is neuroprotective in a mouse model of human retinitis pigmentosa. *Journal of*
400 *Neuroscience*, 34, 8139-8150.

401 Phillips, M. J., Walker, T. A., Choi, H. Y., Faulkner, A. E., Kim, M. K., Sidney, S. S., ... Pardue,
402 M.T. (2008). Tauroursodeoxycholic acid preservation of photoreceptor structure and function in the
403 rd10 mouse through postnatal day 30 . *Investigative Ophthalmology & Visual Science*, 49, 2148–
404 2155.

405 Raghupathy, R. K., Gautier, P., Soares, D. C., Wright, A. F., & Shu, X. (2015) Evolutionary
406 Characterization of the Retinitis Pigmentosa GTPase Regulator Gene. *Investigative Ophthalmology &*
407 *Visual Science*, 56, 6255-6264.

408 Raghupathy, R. K., Zhang, X., Liu, F., Alhasani, R.H., Biswas, L., Akhtar, S., ... Shu, X. (2017)
409 Rpgrip1 is required for rod outer segment development and ciliary protein trafficking in zebrafish.
410 *Scientific Reports*, 7, 16881.

411 Rao, K. N., Li, L., Anand, M., & Khanna, H. (2015) Ablation of retinal ciliopathy
412 protein RPGR results in altered photoreceptor ciliary composition. *Scientific Reports*, 5, 11137.

413 Roche, S. L., Wyse-Jackson, A. C., Byrne, A. M., Ruiz-Lopez, A. M., & Cotter, T. G. (2016)
414 Alterations to retinal architecture prior to photoreceptor loss in a mouse model of retinitis pigmentosa.
415 *The International Journal of Developmental Biology*, 60, 127-139.

416 Roepman, R., Letteboer, S. J., Arts, H. H., van Beersum, S. E., Lu, X., Krieger, E., ... Cremers, F. P.
417 (2005) Interaction of nephrocystin-4 and RPGRIP1 is disrupted by nephronophthisis or Leber
418 congenital amaurosis-associated mutations. *Proceedings of the National Academy of Sciences of the*
419 *United States of America*, 102, 18520-18525.

420 Roque, R. S., Imperial, C. J., & Caldwell, R. B. (1996) Microglial cells invade the outer retina as
 421 photoreceptors degenerate in Royal College of Surgeons rats. *Investigative Ophthalmology & Visual*
 422 *Science*, 37, 196-203.
 423 Schroder, K., & Tschopp, J. (2010) The Inflammasomes. *Cell*, 140, 821–832.
 424 Shu, X., Black, G. C., Rice, J. M., Hart-Holden, N., Jones, A., O'Grady, A., ... Wright, A. F. (2007)
 425 RPGR mutation analysis and disease: an update. *Human Mutation*, 28, 322-328.
 426 Shu, X., Fry, A. M., Tulloch, B., Manson, F. D., Crabb, J. W., Khanna, H., ... Wright, A. F. (2005)
 427 RPGR ORF15 isoform co-localizes with RPGRIP1 at centrioles and basal bodies and interacts with
 428 nucleophosmin. *Human Molecular Genetics*, 14, 1183-1197.
 429 Shu, X., Zeng, Z., Eckmiller, M. S., Gautier, P., Vlachantoni, D., Manson, F. D., ... Wright, A. F.
 430 (2006) Developmental and tissue expression of *Xenopus laevis* RPGR. *Investigative Ophthalmology*
 431 *& Visual Science*, 47, 348-356.
 432 Shu, X., Zeng, Z., Gautier, P., Lennon, A., Gakovic, M., Patton, E. E., Wright, A. F. (2010)
 433 Zebrafish *Rpgr* is required for normal retinal development and plays a role in dynein-based retrograde
 434 transport processes. *Human Molecular Genetics*, 19, 657-670.
 435 Thompson, D. A., Khan, N.W., Othman, M. I., Chang, B., Jia, L., Grahek, G., ... Heckenlively, J. R.
 436 (2012) Rd9 is a naturally occurring mouse model of a common form of retinitis pigmentosa caused by
 437 mutations in RPGR-ORF15. *PLoS One*, 7, e35865.
 438 Tohari, A., Zhou, X., & Shu, X. (2016) Protection against oxidative stress by vitamin D in cone cells.
 439 *Cell Biochemistry and Function*, 34, 82-94.
 440 Vervoort, R., Lennon, A., Bird, A. C., Tulloch, B., Axton, R., Miano, M. G., ... Wright, A. F. (2000)
 441 Mutational hot spot within a new RPGR exon in X-linked retinitis pigmentosa. *Nature Genetics*, 25,
 442 462–466.
 443 Won, J., Gifford, E., Smith, R. S., Yi, H., Ferreira, P. A., Hicks, ... Nishina, P. M. (2009) RPGRIP1 is
 444 essential for normal rod photoreceptor outer segment elaboration and morphogenesis. *Human*
 445 *Molecular Genetics*, 18, 4329-4339.

Won, J., Marín de Evsikova, C., Smith, R. S., Hicks, W. L., Edwards, M. M., Longo-Guess, C., ...
Nishina, P. M. (2011) NPHP4 is necessary for normal photoreceptor ribbon synapse maintenance and
outer segment formation, and for sperm development. *Human Molecular Genetics*, 20, 482-496.
Yanguas-Casás, N., Barreda-Manso, M. A., Nieto-Sampedro, M., & Romero-Ramírez, L. (2017)
TUDCA: An Agonist of the Bile Acid Receptor GPBAR1/TGR5 With Anti-Inflammatory
Effects in Microglial Cells. *J Cellular Physiology*, 232, 2231-2245.
Yoshida N, Ikeda Y, Notomi S, Ishikawa K, Murakami Y, Hisatomi T, Enaida H, Ishibashi T.
Laboratory evidence of sustained chronic inflammatory reaction in retinitis pigmentosa.
Ophthalmology. 2013;120(1):e5-12.
Zeiss, C. J., & Johnson, E. A. (2004). Proliferation of microglia, but not photoreceptors, in the outer
nuclear layer of the rd-1 mouse. *Invest Ophthalmol Vis Sci*. 2004;45(3):971-6.
Zeng HY, Zhu XA, Zhang C, Yang LP, Wu LM, Tso MO. Identification of sequential events and
factors associated with microglial activation, migration, and cytotoxicity in retinal degeneration in rd
mice. *Invest Ophthalmol Vis Sci*. 2005 Aug;46(8):2992-9.
Zhang L, Cui X, Jauregui R, Park KS, Justus S, Tsai YT, Duong JK, Hsu CW, Wu WH, Xu CL, Lin
CS, Tsang SH. Genetic Rescue Reverses Microglial Activation in Preclinical Models of Retinitis
Pigmentosa. *Mol Ther*. 2018. pii: S1525-0016(18)30273-9.
Zhang,T., Baehr,W. and Fu,Y. (2012). Chemical chaperone TUDCA preserves cone photoreceptors in
a mouse model of Leber congenital amaurosis. *Investig. Ophthalmol. Vis. Sci.*, **53**, 3349–3356.
Zhao L, Zabel MK, Wang X, Ma W, Shah P, Fariss RN, Qian H, Parkhurst CN, Gan WB, Wong WT.
Microglial phagocytosis of living photoreceptors contributes to inherited retinal degeneration. *EMBO*
Mol Med. 2015;7(9):1179-97.

Figure Legends

Figure 1 Loss of RPGR caused mislocalization of opsins and transducin. Cryosections of retinas at 1month (M) old were stained for rhodopsin (green), red/green opsin (red), blue opsin (red), transducin (red), and nuclei were counterstained with DAPI (blue). Opsins and transducin were localized to outer segments (OSs) in wildtype (WT) mouse retinas. Rhodopsin was mislocalized to inner segment (IS) in *Rpgr* cko retina, while red/green opsin, blue opsin and transducin exhibited mislocalization to both the IS and outer nuclear layer (ONL) in *Rpgr* cko retina. Arrowheads indicate mislocalized rhodopsin, red/green and blue opsins in the respective images.

Figure 2 NPHP4 was absent from connecting cilia in *Rpgr* cko mice. (A) Paraffin sections of mouse retina were stained for NPHP4 (green) and connecting cilium (CC) marker acetylated α tubulin (red). NPHP4 was localized to connecting cilia and partially co-localized with acetylated α tubulin in wildtype (WT) retina but was absent from *Rpgr* cko (RPGR KO) mouse retinas (n=3). (B) Western blotting with total retina lysis of three WT and *Rpgr* cko mice was performed. (C) NPHP4 (158 kDa) protein level was significantly decreased in *Rpgr* cko mouse retinas at 3, 6 and 12 months (M) old. Data are from two separate Western blots. Statistical comparisons were performed by a non-parametric *t*-test followed by Wilcoxon matched-pairs signed rank test. * $p < 0.05$; ns, no significance.

Figure 3 Cell death occurred in *Rpgr* cko retinas. (A) TUNEL assay was carried out with cryosections of WT and *Rpgr* cko retinas at 1M, 3M, 6M and 12M old. Green signal represented the DNA fragmentation in photoreceptors undergoing cell death. Arrowheads indicate nuclei of apoptotic cells. (B) Quantification of percentage of apoptotic cells in total photoreceptors. Blue line represents wildtype (WT) group and red line represents *Rpgr* cko group. Cell death signal significantly increased at 3M old in cko retinas and peaked at 6m old. Data were collected and analysed by a non-parametric *t*-test followed by Wilcoxon matched-pairs signed rank test (n=5). * $p < 0.05$, ** $p < 0.01$, *** $p < 0.001$; ns, no significance.

Figure 4 Microglia were activated and induced inflammation at an early stage. (A) Immunostaining of microglia marker Iba-1 (red) was performed with cryosections at different ages. In *Rpgr* cko retina, microglia were stimulated into a ramified shape and infiltrated the outer nuclear layer (ONL) as early

as 1M. Arrowheads indicate microglia. (B) The percentage of cells with Iba-1 signal in total photoreceptors in both WT and *Rpgr* cko retina was quantified with ZEN and analysed with a non-parametric *t*-test followed by Wilcoxon matched-pairs signed rank test (n=5). *p<0.05.

Figure 5 Activation of inflammation pathway in photoreceptor cell death. Co-immunostaining was performed for detection of Iba-1 (red) and NLRP3 (green). The overlap of Iba-1 and NLRP3 was observed in the outer nuclear layer (ONL) in cko retina at different ages; in wildtype (WT) retinas, they were found only in outer plexiform layer (OPL). Arrowheads indicate Iba-1/NLRP3.

Figure 6 TUDCA protected the morphological loss of photoreceptors. TUDCA treatment ameliorated retinal degeneration in *Rpgr* cko mice. A total of six eyes (one eye from each treated or untreated mouse) were used for cryosectioning and immunostaining. Nuclei in retina were counterstained with DAPI (blue). (A) Outer nuclear layer (ONL) of untreated *Rpgr* cko retinas (n=6) was significantly thinner than ONL of TUDCA-treated retinas. (B) TUDCA attenuated cell death of photoreceptors as determined by TUNEL assay, indicating that less cell death occurred in ONL after TUDCA treatment. Arrowheads indicate nuclei of apoptotic cells. (C) TUDCA treatment reduced cleaved-caspase 3 protein (red) expression detected by immunostaining. Arrowheads indicate activated caspase 3. Statistical comparisons were performed using one-way ANOVA followed by Bonferroni test. *p<0.05, **p<0.01, ***p<0.001.

Figure 7 TUDCA treatment reduced the activity of microglia in the photoreceptor layer and ameliorated inflammation. A total of six eyes (one eye from each treated or untreated mouse) were used for cryosectioning and immunostaining. (A) Iba-1 (red) represents the microglia state. In the TUDCA treated group, invasion of microglia (arrowheads) in ONL was significantly reduced compared to the untreated group. (B) Co-immunostaining of Iba-1 (red) and NLRP3 (green) was performed. In the TUDCA treated group, NLRP3 expression was restricted to the outer plexiform layer whereas in the untreated group was present also in the ONL. Red arrowheads indicate microglia; green arrowheads indicate NLRP3. Broken lines are used to separate different retinal layers. (C) Immunostaining of IL-1 β (green, arrowheads) was performed and showed that TUDCA treatment dramatically reduced IL-1 β in retinas. Total RNAs were extracted from mouse retinas (one retina

from each eye of individual treated or untreated mice; in total 6 retinas) were used for cDNA synthesis and for qRT-PCR. Statistical comparisons were performed using one-way ANOVA followed by Bonferroni test. * $p < 0.05$, *** $p < 0.001$.

Figure 1

Figure 2

Figure 3

573 **Figure 4**

583 **Figure 5**

597 **Figure 6**

598

599

600

601

602

603

604

605

606

607

608

609

610

611 **Figure 7**

612

613

614

615

616

Supplementary data

Table S1. Primary antibodies were used for this study

Primary Antibody	Dilution	Manufacturer
Rhodopsin	1:200	Abcam, ab98887
M opsin	1:100	Abcam, ab5405
S opsin	1:100	Abcam, ab5407
α -Transducin	1:100	Abcam, ab74059
GRK1	1:100	Abcam, ab2775
NPHP4	1:100	Antibody-online, ABIN1089278
Cleaved Caspase 3	1:100	Cell Signaling #9664
Iba-1	1:100	Wako, 019-19741
NLRP3	1:100	R&D, MAB7578
IL-1 β	1:200	Cell Signaling, #12242
RPGR	1:100	Sigma, HPA001593
Acetylated α tubulin	1:200	Sigma, T6793

Supplementary Figure legends

Figure S1 Full image of Western blot for Figure 2B. Description of the blot is provided in Figure 2B legend.

Figure S2 Morphological change in *Rpgr* cko retinas. Histological examination following hematoxylin and eosin staining showed decreased thickness of the outer nuclear layer (ONL) in *Rpgr* cko retina from 1 month (M) to 12M old. Green bars indicate ONL. Graphs show the thickness of ONL on both superior and inferior sides of retinas. Compared to wildtype (WT, blue line) retinas, the thickness of the ONL of *Rpgr* cko retinas (red line) was significantly decreased from 3months (M) old (n=5). INL, inner nuclear layer; IS, inner segment; OPL, outer plexiform layer; OS, outer segment. Statistical comparisons were performed by a non-parametric *t*-test following by Wilcoxon matched-pairs signed rank test. * $p < 0.05$.

Figure S3 The rhodopsin signal (green) indicated the length of outer segment (OS) in retinas, represented by the double-headed red arrow. IS, inner segment; ONL, outer nuclear layer. (B) The length of outer segments (OS) of *Rpgr* cko (RPGR KO) retinas from 3 months (M) old was significantly reduced when compared to that of wildtype (WT) retinas. The length of OS at different ages was measured by ZEN and analysed by GraphPad Prism (n=5). Statistical comparisons were performed by a non-parametric *t*-test following by Wilcoxon matched-pairs signed rank test. * $p < 0.05$.

Figure S4 Caspase-dependent apoptosis was involved in photoreceptor degeneration in *Rpgr* cko retina. (A) Cryosections of retinas were stained for cleaved-caspase 3 (green), counterstaining with DAPI (blue). Caspase 3 was observed in the outer plexiform layer (OPL) of cko retinas from 3 months (M) old and also in the outer nuclear layer (ONL) at 6M and 12M old. In wildtype (WT) retina, no cleaved-caspase 3 was observed (n=5). (B) Quantification of cleaved-caspase 3 signal. Percentage of photoreceptors in which cleaved-caspase 3 signals were detected in total photoreceptors. Statistical comparisons were performed by a non-parametric *t*-test followed by Wilcoxon matched-pairs signed rank test. * $p < 0.05$.

Figure S5 Counting of photoreceptors in wildtype (WT), TUDCA-treated and untreated mouse retinas. Cryosections of WT, TUDCA-treated and untreated mouse eyes were stained with DAPI (blue). The results represent data obtained from two regions (one in the superior retina and one in the

inferior retinas, 100μm×100μm under 400x magnification, 0.4mm from optic nerve head, ONH) of two sections of each eye. Five eyes from five individual mice were used for the quantification. Data were analyzed by GraphPad Prism with one-way ANOVA.

Figure S6 Expression of Caspase 3 and IL-1β (mRNA) was suppressed by TUDCA treatment.

TUDCA treatment significantly reduced the mRNA levels of Caspase 3 (A) and IL-1β (B) in retinas of *Rpgr* cko mice. Statistical comparisons were performed using one-way ANOVA followed by Bonferroni test. **p<0.01, ***p<0.001.

Figure S1

Figure S3

722 **Figure S4**

723

724

725 **Figure S5**

726

A

B

728