

Hepatitis C reinfection following treatment induced viral clearance among people who have injected drugs

Weir, Amanda; McLeod, Allan; Innes, Hamish; Valerio, Heather; Aspinall, Esther; Goldberg, David; Barclay, Stephen; Dillon, John; Fox, Ray; Fraser, Andrew; Hayes, Peter; Kennedy, Nicholas; Mills, Peter; Stanley, Adrian; Aitken, Celia; Gunson, Rory; Templeton, Kate; Hunt, Alison; McIntyre, Paul; Hutchinson, Sharon

Published in:
Drug and Alcohol Dependence

DOI:
[10.1016/j.drugalcdep.2016.05.012](https://doi.org/10.1016/j.drugalcdep.2016.05.012)

Publication date:
2016

[Link to publication in ResearchOnline](#)

Citation for published version (Harvard):

Weir, A, McLeod, A, Innes, H, Valerio, H, Aspinall, E, Goldberg, D, Barclay, S, Dillon, J, Fox, R, Fraser, A, Hayes, P, Kennedy, N, Mills, P, Stanley, A, Aitken, C, Gunson, R, Templeton, K, Hunt, A, McIntyre, P & Hutchinson, S 2016, 'Hepatitis C reinfection following treatment induced viral clearance among people who have injected drugs', *Drug and Alcohol Dependence*, vol. 165, pp. 53–60.
<https://doi.org/10.1016/j.drugalcdep.2016.05.012>

General rights

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

Take down policy

If you believe that this document breaches copyright please view our takedown policy at <https://edshare.gcu.ac.uk/id/eprint/5179> for details of how to contact us.

Figure 1. Schematic diagram of study cohort.




Figure 2. Frequency of HCV RNA testing during study follow-up. Total number of PWID in each year of study follow-up is given in brackets (excluding those who died during or prior to that year).

