

## **Coenzyme Q10 as a therapeutic candidate for treating inherited photoreceptor degeneration**

Zhang, Xun; Biswas, Lincoln; Tohari, Ali Mohammad; Reilly, James; Tiano, Luca; Shu, Xinhua

*Published in:*  
Neural Regeneration Research

*DOI:*  
[10.4103/1673-5374.221152](https://doi.org/10.4103/1673-5374.221152)

*Publication date:*  
2017

*Document Version*  
Author accepted manuscript

[Link to publication in ResearchOnline](#)

### *Citation for published version (Harvard):*

Zhang, X, Biswas, L, Tohari, AM, Reilly, J, Tiano, L & Shu, X 2017, 'Coenzyme Q10 as a therapeutic candidate for treating inherited photoreceptor degeneration', *Neural Regeneration Research*, vol. 12, no. 12, pp. 1979-1981. <https://doi.org/10.4103/1673-5374.221152>

### **General rights**

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

### **Take down policy**

If you believe that this document breaches copyright please view our takedown policy at <https://edshare.gcu.ac.uk/id/eprint/5179> for details of how to contact us.

1 **Coenzyme Q10 as a potential therapeutic candidate for treating inherited photoreceptor**  
2 **degeneration**

3 Xun Zhang<sup>1</sup>, Lincoln Biswas<sup>1</sup>, Ali Mohammad Tohari<sup>1</sup>, James Reilly<sup>1</sup>, Luca Tiano<sup>2</sup>, Xinhua Shu<sup>1\*</sup>

4 1 Department of Life Sciences, Glasgow Caledonian University, Glasgow G4 0BA, UK

5 2 Department of Life and Environmental Sciences, Polytechnic University of Marche, Ancona 60131,  
6 Italy

7 **Inherited photoreceptor degeneration**

8 The human retina is a highly specialised tissue that enables the perception of light across a range  
9 of intensities and colours. It covers about 65% of the inner surface of the eye and contains three  
10 layers of cells: the outer nuclear layer (ONL) containing the cell bodies and nuclei of the light-  
11 sensitive rod and cone photoreceptors whose photopigment-containing outer segments form the  
12 photoreceptor layer; the inner nuclear layer (INL) containing bipolar, horizontal and amacrine cells;  
13 and the ganglion cell layer (GCL) from which the the optic nerve arises. There are two layers of  
14 synaptic connections between these three layers: the photoreceptors synapse with second order  
15 neurons, mainly bipolar cells, in the outer plexiform layer (OPL), while in turn the bipolar cells form  
16 connections in the inner plexiform layer (IPL) with ganglion cells. The retinal pigment epithelium  
17 (RPE) lies directly behind the photoreceptor layer, is heavily pigmented to reduce scattering of light,  
18 and is essential for the nourishment, maintenance and metabolism of photoreceptors.

19 Inherited photoreceptor degeneration (IPD), usually referred to as monogenic retinal disease, is  
20 characterised by the dysfunction or death of photoreceptor cells (Bramall et al., 2010). IPD has a  
21 large degree of genotypic and phenotypic heterogeneity. Currently, 261 genes and 298 loci have  
22 been reported to be associated with IPD (Retinal Information Network, accessed 16 November 2017,  
23 <<https://sph.uth.edu/RetNet/>>). IPD associated genes function in a wide range of biological  
24 processes, including photoreceptor cell development and maintenance, phototransduction, ciliary  
25 trafficking, phagocytosis, protein quality control, and metabolism (Bramall et al., 2010). Clinical  
26 features in IPD can be classified into rod dystrophy, cone dystrophy, and rod-and-cone dystrophy,

1 depending on which type of photoreceptor cells are initially affected. Photoreceptor cell death in IPD  
2 is mainly through caspase-dependent and caspase-independent apoptosis, though necrosis is also  
3 reported to be involved (Bramall et al., 2010). Both oxidative stress and inflammation play a critical  
4 role in the death of photoreceptor cells.

5

## 6 **Oxidative stress in inherited photoreceptor degeneration**

7 Oxidative stress reflects an imbalance between the production of free radicals and their  
8 counteraction by the antioxidant defence system (Wright et al., 2004). Photoreceptor cells consume  
9 oxygen at possibly the highest rate in the body, producing extremely large quantities of reactive  
10 oxygen and nitrogen species (RONS) and subsequently causing oxidative damage to macromolecules  
11 such as nucleic acids, proteins and lipids (Zhang et al., 2017). Previously we investigated oxidative  
12 stress in the retinae of four IPD mouse models (*Rho*<sup>-/-</sup>, *Prph2*<sup>rd5/rd5</sup>, *Pde6b*<sup>rd1/rd1</sup> and *Pde6b*<sup>atrd1/atrd1</sup>) and  
13 found that they had significantly decreased activities of mitochondrial complex I compared to that of  
14 wildtype retinae, an effect that would have been compounded by the consequent increased  
15 production of reactive oxygen species (ROS) by the mitochondria. *Rho*<sup>-/-</sup> and *Pde6b*<sup>rd1/rd1</sup> mouse  
16 retinae also had markedly lower levels of the reduced form of the antioxidant glutathione (GSH)  
17 (Vlachantoni et al., 2011). Patients with retinitis pigmentosa, a common IPD, exhibited ocular  
18 oxidative damage with significantly reduced total antioxidant capacity and a reduced ratio of GSH to  
19 GSSG (the oxidised form of glutathione) in the aqueous humour compared to that of controls  
20 (Campochiaro et al., 2015). A mix of antioxidants ( $\alpha$ -tocopherol, ascorbic acid,  $\alpha$ -lipoic acid and  
21 Mn(III)tetrakis (4-benzoic acid) porphyrin) significantly reduced cone death and preserved cone  
22 function in an IPD mouse model (rd1), possibly through inhibiting oxidative damage (Komeima et al.,  
23 2006). Co-expression of catalase and SOD2 in the photoreceptor cells of an IPD mouse model (rd10)  
24 significantly reduced superoxide radicals and carbonyl contents, decreased cone cell death, and  
25 preserved cone function (Usui et al., 2009).

1 Light exposure also generates RONS and this makes photoreceptor cells vulnerable to oxidative  
2 damage. Accumulated evidence demonstrates that light damage to photoreceptors is due to the  
3 accumulation of all-trans retinal released from photo-activated rhodopsin, particularly when the  
4 recycling of all-trans retinal to 11-cis retinal is blocked due to genetic mutations. All-trans retinal  
5 molecules can form toxic bis-retinoids, such as N-retinylidene-N—retinylethanolamine (A2E), which  
6 accumulate in the RPE cells through phagocytosis (Organisciak & Vaughan, 2010). Most components  
7 of bis-retinoids are able to produce RONS when undergoing photo-oxidation. Synthetic antioxidants  
8 such as dimethylthiourea have been shown to protect photoreceptors from light-induced damage,  
9 functioning as a quencher of free radicals. Natural products also demonstrate a protective role  
10 against light-induced retinal damage: some substances act directly as antioxidants, while others are  
11 involved in up-regulation of antioxidant gene expression.

12

### 13 **Microglia I activation in inherited photoreceptor degeneration**

14 Microglia are a type of macrophage resident in the nervous system including the brain and retina.  
15 In the developing retina, microglia precursors may move from retinal vascular vessels and the ciliary  
16 body into the retina. After entering the retina, these precursors move along the axon fascicles to  
17 their final location in the retina. Microglia ultimately settle in the plexiform layers of the retina  
18 where their development completes. They exhibit a ramified form, constantly moving to surveille  
19 the retinal neurons. The cellular markers and receptors of microglia indicate the status of microglia  
20 (i.e. inactive or active form). These cell surface receptors also provide microglia with special immune  
21 functions. Microglia have a phagocytic role in engulfing presynaptic inputs. This function depends on  
22 microglia-specific phagocytic signalling pathway, complement receptor 3 (CR3)/C3. Under healthy  
23 conditions, microglia fulfil specialized immune tasks by releasing cytokines and neurotrophic factors.  
24 A typical interaction between microglia and neurons is the FKN/CX2CR1 system (fractalkine or  
25 CX3CL1 and its G protein- coupled receptor, CX3CR1). CX3CL1 is a chemokine in the form of a  
26 membrane-bound protein, which can bind to CX3CR1 on microglia and which modulates the

1 activation of microglia (Vecino et al., 2016). CX3CR1 signalling in retinal microglia plays an important  
2 role in controlling motility of the thin microglial process that continuously monitor the  
3 microenvironment in the healthy retina. Following retinal damage or a change in its condition (such  
4 as inflammation, light injury or genetic defects), microglia become activated and change their  
5 morphology, with expression of surface markers such as complement receptor, MHC-II, and CD68.  
6 Subsequently, activated microglia exhibit amoeboid movement and enter the photoreceptor layer to  
7 carry out phagocytosis (Karlstetter et al., 2015).

8       Activated microglia may play a pathological role in the degenerating photoreceptor layer by  
9 releasing neurotoxic factors, such as TNF- $\alpha$  and IL-1 $\beta$ . In IPD, evidence that the dying rods may  
10 attract the resting microglia to mediate phagocytosis in retinitis pigmentosa patients was first  
11 presented by Gupta. (2003). Activated microglia were found in the region of dying rods and it was  
12 proposed that the initial rod cell death further triggers the onset of cone death. In the widely used  
13 rd1 mouse model which contains nonsense mutation in PDE6b gene, activated microglia were  
14 detected from postnatal day 12 in the outer nuclear layer with releasing cytokines (such as TNF- $\alpha$ )  
15 and chemokines C-C motif ligand 2 (CCL2), C-C motif ligand 5 (CCL5) and monocyte chemoattractant  
16 protein 1 (MCP1) (Karlstetter et al., 2015). Similarly, in rd10 mouse, the rod degeneration is  
17 accompanied by infiltration of microglia. The dying rods exhibited the “eat me” signal PtdSer on the  
18 cell surface to induce the invasion of activated microglia into the ONL. Genetic ablation of retinal  
19 microglia significantly reduced rod cell death in rd10 mice (Zhao et al., 2015). Inhibition of microglia  
20 activation by minocycline also protected photoreceptor cells from apoptosis in IPD mouse models  
21 rds and rd10 (Bramall et al., 2010; Peng et al., 2014). Given this evidence, activated microglia-  
22 mediated phagocytosis appears to be the underlying cell death mechanism contributing to retinal  
23 degeneration in IPD.

24

25 **Coenzyme Q10 protects retinal cells from oxidative stress and inhibits retinal microglia activation**

1 Coenzyme Q10 (CoQ10) is a vitamin-like molecule that is endogenously synthesized in all aerobic  
2 organisms, existing as oxidized (ubiquinone), partially reduced (semiquinone) and reduced form  
3 (ubiquinol) within the body. The lipophilic molecule plays a central role in cellular growth and  
4 metabolism through the electron transport chain where it functions as an electron acceptor,  
5 collecting electrons from mitochondrial respiratory chain complex I and complex II, then shuttling  
6 them to complex III. Additionally, CoQ10 plays a role as an antioxidant, acting as a free radical  
7 scavenger and so protecting cells from oxidative damage. It is particularly abundant in tissues that  
8 are highly oxygen-demanding, such as cardiac muscle. Although present in the retina, levels are  
9 comparatively low relative to other tissues dependent on oxidative metabolism. Following a peak in  
10 the third decade of life, the level of CoQ10 declines with age due to decreased synthesis and  
11 increased degradation. CoQ10 supplementation has been shown to slow the progression of age-  
12 related neurodegenerative diseases, such as Parkinson's disease and age-related macular  
13 degeneration (Zhang et al., 2017).

14 Previous work demonstrated that CoQ10 treatment protected RGC-5 cells (a rat ganglion cell  
15 line) from oxidative damage caused by exposure to H<sub>2</sub>O<sub>2</sub>, antimycin (mitochondria complex III  
16 inhibitor), radiation, or serum starvation (Zhang et al., 2017). We have also treated human retinal  
17 pigment epithelium cells (ARPE-19) with H<sub>2</sub>O<sub>2</sub> only or with H<sub>2</sub>O<sub>2</sub> and CoQ10, and found that cell  
18 viability was significantly increased in ARPE-19 cell exposure to CoQ10 and H<sub>2</sub>O<sub>2</sub> when compared to  
19 cells exposed to H<sub>2</sub>O<sub>2</sub> alone (data not shown). In addition, inflammatory cytokines such as IL-1 $\beta$  were  
20 markedly decreased and expression of antioxidant genes (e.g. SOD1) was notably increased in ARPE-  
21 19 cells co-exposed to CoQ10 and H<sub>2</sub>O<sub>2</sub> when compared to cells treated with H<sub>2</sub>O<sub>2</sub> only (data not  
22 shown).

23 Senescence accelerated P8 (SAMP8) mice have high levels of protein oxidation and lipid  
24 peroxidation in examined tissues, demonstrate some pathological features similar to those  
25 presented in age-related neurodegenerative diseases, and have been widely used as a model for  
26 oxidative stress and ageing diseases (Morley, et al., 2012). To investigate the effect of CoQ10 on

1 retinal microglial activation, we treated SAMP8 mice with CoQ10 via oral administration (500mg/kg,  
2 daily) over a two-month period. We found that microglia migrated into the inner portion of the ONL  
3 in untreated SAMP8 mice, while microglia were settled in the OPL in CoQ10-treated SAMP8 mice.  
4 We also observed that expression of inflammatory cytokine IL-1 $\beta$  in the OPL was noticeably  
5 increased in untreated SAMP8 mice when compared to CoQ10-treated mice (Figure 1).

6

## 7 **Future directions**

8 Although CoQ10 supplement treatment shows protective effects in age-related neurodegenerative  
9 diseases and inhibits oxidative damage, inflammation and microglia activation in retina cell lines and  
10 SAMP8 retinae, more pre-clinical studies in IPD animal models are needed before it can be used  
11 effectively in a therapeutic setting. Mutations in the RPGRIP1 gene can cause Leber congenital  
12 amaurosis (LCA), juvenile retinitis pigmentosa (RP) and cone-rod dystrophy (common types of IPD) in  
13 humans. Recently, we characterized a zebrafish IPD model that carries a nonsense mutation in the  
14 RPGRIP1 gene (Raghupathy et al., in press). RPGRIP1 mutant zebrafish did not develop  
15 photoreceptor outer segments and displayed mislocalization of rhodopsin, early rod degeneration  
16 and subsequently cone degeneration (Figure 2) (Raghupathy et al., in press), similar to clinical  
17 features presented in LCA and juvenile RP patients. The potential for delaying photoreceptor cell  
18 death by CoQ10 can be evaluated in this IPD zebrafish. More evaluation may be needed in other  
19 mouse IPD models. Based on the preclinical data, clinical trial can be carried out in IPD patients. In  
20 conclusion, CoQ10 has the capacity to inhibit oxidative stress, decrease inflammation and suppress  
21 microglia activation, which will offer potential therapy for IPD patients.

## 22 **Acknowledgement**

23 Work in Dr Shu's lab was supported by the Rosetrees Trust (M160 and M160-F1), the Fight for Sight,  
24 the Glasgow Children's Hospital Charity (YRSS/PSG/2014) and the Visual Research Trust (VR2014).

## 25 **References**

1 Bramall AN, Wright AF, Jacobson SG and McInnes RR (2010) The genomic, biochemical, and cellular  
2 responses of the retina in inherited photoreceptor degenerations and prospects for the treatment of  
3 these disorders. *Annu Rev Neurosci* 33:441-472.

4 Campochiaro PA, Strauss RW, Lu L, Hafiz G, Wolfson Y, Shah SM, Sophie R, Mir TA and Scholl HP  
5 (2015) Is there excess oxidative stress and damage in eyes of patients with retinitis pigmentosa?  
6 *Antioxid Redox Signal* 23:643-648.

7 Gupta N, Brown KE and Milam AH (2003) Activated microglia in human retinitis pigmentosa, late-  
8 onset retinal degeneration, and age-related macular degeneration. *Exp Eye Res* 76:463-471.

9 Karlstetter M, Scholz R, Rutar M, Wong WT, Provis JM and Langmann T (2015) Retinal microglia: just  
10 bystander or target for therapy? *Prog Retin Eye Res* 45:30-57.

11 Komeima K, Rogers BS, Lu L and Campochiaro PA. Antioxidants reduce cone cell death in a model of  
12 retinitis pigmentosa. *Proc Natl Acad Sci U S A*. 2006 Jul 25;103(30):11300-5

13 Morley JE, Armbrecht HJ, Farr SA and Kumar, VB (2012) The senescence accelerated mouse (SAMP8)  
14 as a model for oxidative stress and Alzheimer's disease. *Biochem Biophys Acta* 1822 (5): 650-6.

15 Organisciak DT and Vaughan DK (2010) Retinal light damage: mechanisms and protection. *Prog Retin*  
16 *Eye Res* 29:113-134.

17 Peng B, Xiao J, Wang K, So KF, Tipoe GL and Lin B (2014) Suppression of microglial activation is  
18 neuroprotective in a mouse model of human retinitis pigmentosa. *J Neurosci*. 34:8139-8150.

19 Raghupathy RK, Zhang X, Liu L, Alhasani RH, Biswas L, Akhtar S, Pan L, Moens CB, Li W, Liu M,  
20 Kennedy BN and Shu X. *Rpgrp1* is required for rod outer segment development and ciliary protein  
21 trafficking in zebrafish. *Scientific Reports*. In press.

22 Usui S, Komeima K, Lee SY, Jo YJ, Ueno S, Rogers BS, Wu Z, Shen J, Lu L, Oveson BC, Rabinovitch PS  
23 and Campochiaro PA (2009) Increased expression of catalase and superoxide dismutase 2 reduces  
24 cone cell death in retinitis pigmentosa. *Mol Ther* 17:778-786.

25 Vecino E, Rodriguez FD, Ruzafa N, Pereiro X and Sharma SC (2016) Glia-neuron interactions in the  
26 mammalian retina. *Prog Retin Eye Res* 51:1-40.



1 Vlachantoni D, Bramall AN, Murphy MP, Taylor RW, Shu X, Tulloch B, Van Veen T, Turnbull DM,  
2 Wright AF, Jacobson SG, Cideciyan AV, Roman AJ, Shu X, Vlachantoni D, McInnes RR and Riemersma  
3 RA (2004) Lifespan and mitochondrial control of neurodegeneration. *Nat Genet* 36:1153-1158.  
4 Zhang X, Tohari AM, Marcheggiani F, Zhou X, Reilly J, Tiano L and Shu X (2017) Therapeutic potential  
5 of co-enzyme Q10 in retinal diseases. *Curr Med Chem*. doi: 10.2174/0929867324666170801100516.  
6 [Epub ahead of print]  
7 Zhao L, Zabel MK, Wang X, Ma W, Shah P, Fariss RN, Qian H, Parkhurst CN, Gan WB and Wong WT  
8 (2015) Microglial phagocytosis of living photoreceptors contributes to inherited retinal  
9 degeneration. *EMBO Mol Med* 7:1179–1197.

10

11

12

13

14

15

16

17

18

19

20

21

22

23


24

25

26

27

1 **Figures and legends**



2

3 **Figure 1** CoQ10 treatment blocked microglia migration to outer nuclear layer (ONL) and suppressed  
4 inflammation. Untreated and CoQ10-treated SAMP8 mouse eyes were subjected to cryosection and  
5 immunostaining with anti-Iba-1 antibody (labelling microglia) and anti-IL-1β antibody. In untreated  
6 (No Q<sub>10</sub>) SAMP8 mouse retina, microglia (arrowheads in left-hand boxes) migrated into the inner  
7 portion of ONL; in CoQ10 treated (Q<sub>10</sub>) SAMP8 retina, microglia stayed in the outer plexiform layer  
8 (OPL). Expression of inflammatory cytokine IL-1β (arrowheads in right-hand boxes), was noticeably  
9 higher in OPL of untreated SAMP8 retina, but not in CoQ10 treated retina. (INL: inner nuclear layer).


10

11

12

13

14



1

2 **Figure 2** RPGRIP1<sup>-/-</sup> zebrafish displayed mislocalization of rhodopsin, early rod degeneration and  
 3 subsequent cone degeneration. Eyes from both wildtype and RPGRIP1 mutant zebrafish at 14 days  
 4 (14dpf) and 6 months old (6mpf) were subjected to cryosection and immunostaining using 4D2  
 5 antibody (labelling rod outer segments, ROS), mislocalization of rhodopsin was seen in mutant retina  
 6 at 14dpf, while rod cells showed early signs of degeneration. All rod cells in mutant retina were  
 7 degenerated at 6mpf. Zebrafish have long cones (red and green cones) and short cones (blue and  
 8 UV cones), methylene blue staining showed that only short cones remained in the retina of 13mpf  
 9 RPGRIP1 mutant zebrafish. GCL, ganglion cell layer; INL, inner nuclear layer; ONL, outer nuclear  
 10 layer; RPE, retinal pigment epithelium.