

Effectiveness of interventions to improve the public's antimicrobial resistance awareness and behaviours associated with prudent use of antimicrobials: a systematic review

Price, Lesley; Gozdzielewska, Lucyna; Young, Mairi; Smith, Fraser; MacDonald, Jennifer; McParland, Joanna; Williams, Lynn; Langdridge, Darren; Davis, Mark; Flowers, Paul

Published in:
Journal of Antimicrobial Chemotherapy

DOI:
[10.1093/jac/dky076](https://doi.org/10.1093/jac/dky076)

Publication date:
2018

Document Version
Author accepted manuscript

[Link to publication in ResearchOnline](#)

Citation for published version (Harvard):

Price, L, Gozdzielewska, L, Young, M, Smith, F, MacDonald, J, McParland, J, Williams, L, Langdridge, D, Davis, M & Flowers, P 2018, 'Effectiveness of interventions to improve the public's antimicrobial resistance awareness and behaviours associated with prudent use of antimicrobials: a systematic review', *Journal of Antimicrobial Chemotherapy*, vol. 73, no. 6, pp. 1464-1478. <https://doi.org/10.1093/jac/dky076>

General rights

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

Take down policy

If you believe that this document breaches copyright please view our takedown policy at <https://edshare.gcu.ac.uk/id/eprint/5179> for details of how to contact us.

Manuscript category: Systematic review

Title: The effectiveness of interventions to improve the public's antimicrobial resistance awareness and behaviours associated with prudent use of antimicrobials: a systematic review

Running title: Effectiveness of AMR interventions: systematic review

Authors: Lesley PRICE^{1*}, Lucyna GOZDZIELEWSKA¹, Mairi YOUNG¹, Fraser SMITH¹, Jennifer MACDONALD¹, Joanna MCPARLAND¹, Lynn WILLIAMS², Darren LANGDRIDGE³, Mark DAVIS⁴, Paul FLOWERS¹.

¹ Safeguarding Health through Infection Prevention Research Group, Glasgow Caledonian University, Cowcaddens Road, Glasgow, G4 0BA, UK

² School of Psychological Sciences and Health, University of Strathclyde, 40 George Street, Glasgow, G1 1QE

³ Faculty of Arts & Social Sciences, Open University, Walton Hall, Milton Keynes, MK7 6AA

⁴ School of Social Sciences, Monash University, Chancellors Walk, Melbourne, Victoria, Australia, 3800

Corresponding author details: Dr Lesley PRICE, Institute for Applied Health Research, Glasgow Caledonian University, Cowcaddens Road, Glasgow, G4 0BA, UK. L.Price@gcu.ac.uk, Phone +44 (0)141 331 3431. Fax +44 (0)141 331 3005.

1 **Synopsis**

2 **Background:** A global antimicrobial resistance (AMR) awareness intervention targeting the
3 general public has been prioritised.

4 **Objectives:** To evaluate the effectiveness of interventions which aim to change AMR
5 awareness and subsequent stewardship behaviours amongst the public.

6 **Methods:** Five databases were searched between 2000 and 2016 for interventions to change
7 the public's AMR awareness and/or antimicrobial stewardship behaviours. Study designs
8 meeting the EPOC criteria: non-controlled before and after studies and prospective cohort
9 studies were considered eligible. Participants recruited from healthcare settings and studies
10 measuring stewardship behaviours of healthcare professionals were excluded. Quality of
11 studies was assessed using EPOC risk of bias criteria. Data were extracted and synthesised
12 narratively.

13 **Results:** Twenty studies were included in the review with nine meeting the EPOC criteria. The
14 overall risk of bias was high. Nineteen studies were conducted in high-income countries. Mass
15 media interventions were most common (n = 7), followed by school-based (n = 6) and printed
16 materials interventions (n = 6). Seventeen studies demonstrated a significant effect on
17 changing knowledge, attitudes, or the public's antimicrobial stewardship behaviours. Analysis
18 showed that interventions targeting schoolchildren and parents have a notable potential but
19 for the general public the picture is less clear.

20 **Conclusions:** Our work provides an in-depth examination of the effectiveness of AMR
21 interventions for the public. However, the studies were heterogeneous and the quality of

22 evidence was poor. Well-designed, experimental studies on behavioural outcomes of such
23 interventions are required.

24 **Registration:** PROSPERO International prospective register of systematic reviews (PROSPERO
25 2016:CRD42016050343).

26

27 **Introduction**

28 The rise of antimicrobial resistance (AMR) is a rapidly developing global threat that greatly affects
29 our ability to deliver effective healthcare and results in a financial burden.¹ AMR refers to the ability
30 of a microorganism to adapt and grow despite the presence of antimicrobials. AMR threatens
31 effective treatment of an ever-increasing range of infections.¹ Therefore, increasing AMR is
32 becoming a major public health concern. Although AMR is a naturally occurring phenomenon,
33 inappropriate use of antimicrobials is the main driver of AMR.¹ The demands for the use of
34 antimicrobials are increasing worldwide and because of suboptimal management of these demands,
35 huge quantities of antimicrobials are being misused.² Together these highlight the need for effective
36 strategies encouraging prudent use of antimicrobials.

37 The O'Neil report emphasises the need for AMR awareness interventions directed towards the
38 public and development of a uniform, globally consistent set of AMR messages that could be then
39 tailored to meet the specific demands of local settings.² However, the report does not provide
40 recommendations on components of such interventions.²

41 Previous evidence syntheses shows that the overall levels of knowledge and understanding of AMR
42 amongst the public is generally low and members of the public often lack an understanding of their
43 potential contribution to the development of AMR.³⁻⁵ Although high-level evidence demonstrating
44 the effectiveness of interventions in increasing public understanding of AMR exists,⁵⁻⁷ these
45 evaluations are methodologically diverse. It is therefore challenging to identify what interventions
46 work, why and for whom in order to inform future interventions.

47 Thus, the aim of this systematic review is to provide the best quality evidence regarding the
48 effectiveness of AMR interventions that change public awareness and their subsequent antimicrobial
49 stewardship behaviours. Although, antimicrobial stewardship is most commonly thought of in
50 medical settings, the word "stewardship" means "taking care of" particularly on behalf of others.

51 Furthermore, a One Health perspective requires the *collaborative* effort of all stakeholders to take
52 the responsibility for the prudent use of antimicrobials. Therefore, within this work, we use the term
53 “antimicrobial stewardship” to explore the public’s behaviours related to their prudent use of
54 antimicrobials (such as, but not limited to: adhering to prescribers’ directions, not taking or
55 demanding antimicrobial prescription for colds and flu and safe disposal of leftover antimicrobials).
56 We believe an understanding of the public’s antimicrobial stewardship is central to engaging them
57 with their part to play in reducing the drivers of AMR on behalf of future generations, other key
58 stakeholders such as prescribers, and the global community.

59 **Methods**

60 This review was prospectively registered on the PROSPERO International prospective register of
61 systematic reviews (PROSPERO 2016:CRD42016050343 Available from:
62 http://www.crd.york.ac.uk/PROSPERO/display_record.asp?ID=CRD42016050343), and is reported in
63 accordance with the Preferred Reporting Items for Systematic Reviews and Meta-Analyses (PRISMA)
64 statement.⁸

65 ***Search strategy***

66 CINAHL, Cochrane Library, EMBASE, MEDLINE and PsycINFO databases were searched for articles
67 published between 2000 and 2016 using keywords associated with the following four concept areas:
68 (1) population - general public; (2) intervention - interventions designed to increase antimicrobial
69 awareness and/or to improve antimicrobial stewardship behaviour amongst the general public; (3)
70 context - AMR or the public’s antimicrobial stewardship; outcomes - all relevant short, medium or
71 long-term outcomes related to the public’s antimicrobial resistance and/or antimicrobial
72 stewardship behaviours (knowledge/awareness, learning, public behavioural and cognition
73 outcomes). The search strategy incorporated controlled vocabulary thesaurus terms and free text
74 words contained in titles and abstracts. No restrictions were applied to language and publication

75 status. The search strategy was amended according to the functionality of each of the databases. An
76 example of the search strategy applied to MEDLINE is presented in Table S1 (available at JAC Online).
77 In addition to the database search, the reference lists of included papers and previous systematic
78 reviews were searched manually and citation searches were conducted through Web of Science in
79 order to identify additional records.

80 ***Study selection***

81 Cochrane's Effective Practice and Organization of Care (EPOC) recommendations⁹ were used to
82 initially select studies for inclusion in the review. Although, EPOC guidelines suggest the inclusion of
83 Randomised controlled trials (RCT), Non-randomised controlled trials(NRT), controlled before and
84 after studies (CBA), Interrupted time series studies (ITS) and repeated measures studies exclusively,⁹
85 because of the limited number of eligible studies meeting the EPOC criteria, non-controlled before
86 and after studies and prospective cohort studies were also deemed eligible to the review.
87 Interventions targeting the general public population and designed to increase public antimicrobial
88 awareness and/or to improve the public's antimicrobial stewardship were deemed eligible to the
89 review. Furthermore, time bound geographical controls or no exposure comparators and all relevant
90 short, medium or long-term outcomes related to antimicrobial resistance and/or the public's
91 antimicrobial stewardship behaviours were included, while those related to antimicrobial prescribing
92 were excluded as this was regarded as the behaviour of healthcare professionals rather than of the
93 members of the general public. Eligibility criteria applied in this study are presented in detail in Table
94 1.

95 Titles and abstracts of identified records were screened against the eligibility criteria (Table 1) by
96 one of three reviewers (MY, LG, FS) with a 30% sub-set of excluded studies independently checked
97 by another reviewer (MY, LG or FS). The level of agreement on this sub-set was 99%. Disagreements
98 were resolved with the involvement of another, experienced reviewer (LP). Full-texts of papers
99 which appeared to meet the inclusion criteria, or where there was insufficient information within

100 the title and abstract were screened by two out of three independent reviewers (LG, MY, FS), with a
101 fourth reviewer (LP) checking all decisions and resolving any discrepancies. Whenever possible,
102 foreign-language papers were translated by members of the team who have a command in foreign
103 languages (LG, JP), or were translated using Google Translate.

104 ***Data extraction and quality assessment***

105 Two out of three reviewers (LG, MY, FS) independently extracted data from eligible studies using
106 standardised tool, designed for the purpose of the study (Table S2, available at JAC online).

107 For studies that met the EPOC study design criteria (RT, NRT, CBA, ITS),⁹ risk of bias was assessed
108 across domains by one reviewer (MY, LG or FS) and checked by a second reviewer (MY, LG or FS)
109 using standard EPOC risk of bias criteria.⁹ Disagreements were resolved through consensus or, if
110 necessary consulted with fourth reviewer (LP). Risk of bias assessments were not conducted for non-
111 controlled before and after studies as it was assumed that the risk of bias of these studies was high.
112 No studies were excluded based on quality assessment.

113 ***Data analysis***

114 Given the heterogeneity of the study designs, populations, interventions and outcome measures, it
115 was not possible to pool the results in a meta-analysis. Therefore, we applied an alternative,
116 systematic approach to assessing complex interventions and carried out a narrative synthesis of
117 evidence following the Cochrane Consumers and Communication Review Group's guidelines.¹⁰
118 Individual study characteristics and findings were summarised and similarities, differences and
119 patterns identified. Studies were grouped to those meeting or not meeting the EPOC criteria and
120 categorised according to the target population. To identify discernible patterns of effectiveness,
121 identified studies were mapped across five categories of intervention effectiveness. These categories
122 were based upon both the strength of the evidence and the position of the primary outcome within
123 the casual chain linking antecedents of behaviour to actual behaviour change. For example,

124 knowledge is understood to be a necessary but insufficient predictor of behaviour as people can
125 develop good awareness and understanding of AMR yet still fail to implement the public's AMR
126 stewardship. The five categories of a relative measure of effectiveness included (1) interventions
127 indicative of clear positive behaviour change in the desired direction, (2) interventions indicative of
128 some positive behaviour change in the desired direction, (3) interventions indicative of positive
129 effect on the antecedent of behaviour, such as knowledge or awareness in the desired direction, (4)
130 interventions indicative of no effect on behaviour or antecedents of behaviour, and (5)
131 interventions indicative of negative effect on behaviour or antecedents of behaviour in a non-
132 desired direction.

133 **Results**

134 Electronic search resulted in the total of 17,312 records. An additional 31 records were identified
135 through reference and citation searching of the included papers. A total number of 60 studies that
136 did not meet eligibility criteria were excluded during the full text reviewing stage. Articles were
137 excluded for not meeting study design criteria, study participants being recruited from healthcare
138 settings, context other than AMR, study outcomes not related to the public's AMR awareness or
139 antimicrobial stewardship, full text record being unavailable, and other reasons such as record being
140 a study protocol, conference abstract of already identified study, short report of already identified
141 study, inability to translate non-English paper, or majority of participants recruited for the study
142 being healthcare workers. A detailed list of excluded papers is presented in Table S3 (available at JAC
143 Online). Following screening, 20 studies that matched the eligibility criteria were included in the
144 review. A detailed process of study selection is presented in Figure 1.

145 **Study characteristics**

146 As shown in Table 2, study designs of the 20 reviewed studies included randomised controlled trials
147 (n = 2),^{11, 12} non-randomised controlled trials (n = 3),¹³⁻¹⁵ controlled before-after studies (n = 4),¹⁶⁻¹⁹
148 non-controlled before-after studies (n = 10)²⁰⁻²⁹ and a prospective cohort study (n = 1).³⁰

149 Apart from one study conducted in Moldova,¹⁶ all studies were conducted in high-income countries,
150 with the majority conducted in the United States (US) (n = 8),^{11, 12, 14, 15, 19, 23, 28, 30} or in the United
151 Kingdom (UK) (n = 5).^{18, 22, 24, 26, 27} The remaining studies were conducted in Italy,¹³ Portugal,²⁰
152 Poland,²⁵ New Zealand,²¹ and Australia,²⁹ while one study was a multisite study conducted in the UK,
153 Czech Republic and France¹⁷ (Table 2).

154 As shown in Table 2, the most common types of interventions were mass media interventions.^{13, 14,}
155 ^{18, 24, 25, 29, 30} Apart from Madle *et al.* (2008)²⁴ who used website only, all mass media interventions
156 were multimodal and used a variety of outlets, such as billboards, radio, television, newspapers,
157 magazines, websites, and printed resources such as, posters, brochures, pamphlets, leaflets, stickers
158 or badges distributed to community sites or healthcare settings. Six studies used printed materials
159 interventions, either alone,^{12, 15, 21, 23} or in combination with educational presentations,^{11, 19} while
160 Stockwell *et al.* (2010)²⁸ delivered taught modules to Latino community parents. Other interventions
161 were school based, and included a student peer-taught program,¹⁶ e-bug web game,²² interactive
162 workshops,²⁷ school lessons delivered using the “Bug Investigators” pack²⁶ and presentation
163 followed by discussion,²⁰ while the intervention delivered in the study by Lecky *et al.* (2010)¹⁷
164 involved the delivery of a lesson, printed materials, interactive activities and question & answer
165 session.

166 Comparators were similar across the ten controlled studies.^{11-19, 30} With an exception of Lecky *et al.*
167 (2010)¹⁷ who compared the educational intervention to the usual school curriculum, control groups
168 were not exposed to the interventions.

169 Table 2 shows that the most common outcome measure was change in knowledge, attitudes or
170 beliefs, measured alone (n = 10)^{11-13, 15, 17, 20, 22, 24, 26, 27} or in combination with change in the public's
171 antimicrobial stewardship behaviour (n = 8).^{16, 18, 19, 21, 23, 25, 28, 29} Two studies measured the public's
172 antimicrobial stewardship behaviour outcomes exclusively (n = 2).^{14, 30}

173 ***Quality of studies***

174 Amongst the included studies, nine met the EPOC study design criteria.⁹ As shown in Table 3, the
175 overall risk of bias of the included studies was generally high. Apart from one study,¹¹ all had at least
176 one item assessed as high risk with the number of high risk items ranging from one¹⁵ to five.¹⁸ High
177 risk of bias was most commonly associated with generation of sequence allocation, risk of
178 contamination and other risks.

179 For all studies that met the EPOC criteria, insufficient information was provided for at least 2 of the
180 items which were regarded as unclear risk. None of the studies provided information regarding
181 blinding, and in all studies reporting secondary outcomes, the risk of bias for incomplete secondary
182 outcome data could not be assessed.¹³⁻¹⁶ The number of low risk items ranged from one^{17, 18} to five
183 ¹⁶ with the "selective outcome reporting" item being most commonly assessed as low risk. None of
184 the studies had a low risk score for any of the following items: "allocation concealment", "addressing
185 incomplete secondary outcome data" and "blinding primary outcome data".

186 Risk of bias was not assessed for the non-controlled before-after studies²⁰⁻²⁹ and a prospective
187 cohort study.³⁰ These study designs did not meet the EPOC criteria;⁹ therefore, it was assumed that
188 the risk of bias of these studies was high.

189 ***Relative effectiveness of interventions***

190 Reviewed interventions were grouped into five categories of relative measure of effectiveness. As
191 shown in Table 4, six studies demonstrated a clear desired behaviour change following the
192 intervention, while two studies resulted in some desired behaviour change. Desired effect on the
193 antecedent of behaviour was reported in nine papers. One study showed no effect, while two
194 studies demonstrated an increase in drivers of AMR following the intervention.

195 ***Effectiveness of interventions delivered to populations through the lifecycle***

196 In 17 of the studies, the intervention had a significant effect on the outcome of interest amongst the
197 populations through the lifecycle. These included schoolchildren, university students, parents and
198 the general public.

199 ***Schoolchildren***

200 All six school-based educational interventions that targeted schoolchildren aged between 9-15 years
201 ^{16, 17, 20, 22, 26, 27} found a significant increase in knowledge following the educational intervention (Table
202 2). However, Farrell *et al.* (2011)²² found a significant knowledge change in only 3 out of 21
203 questions ($p \leq 0.02$), and no overall change in knowledge. The three questions for which significant
204 improvement was reported related to the valuableness of “good microbes”, the presence of
205 microbes despite inability to see them, and handwashing being an effective method of removing
206 microbes from the hands. Only one study ¹⁶ measured behavioural outcome in addition to beliefs,
207 and found that children in the intervention group were 3.2 times more likely than other students to
208 report that they had not taken an antibiotic for a cold or flu ($p < 0.001$).

209 None of the studies measured long term outcomes of school-based interventions. Post-intervention
210 outcomes were measured immediately following the intervention,²² or between 1-8 weeks after the
211 intervention. In addition, one study ¹⁷ found that the increase in knowledge was maintained at 6
212 week post intervention in junior but not for senior school students.

213 *University Students*

214 University students were targeted in one experimental study that aimed to investigate whether an
215 educational intervention (information booklet) resulted in an increase of young adult consumers'
216 preference for physicians who do not unnecessarily prescribe antibiotics for simple acute upper
217 respiratory tract infections.¹⁵ This study demonstrated, that exposure to the intervention
218 significantly increased the mean preferred start date for antibiotics after the onset of an infection
219 from 2.3.-3.9 days ($p < 0.1$) and preference for a physician who would not prescribe antibiotics at day
220 3 of an infection ($p < 0.1$). However, this was still well before recommended time of 10-14 days.

221 *Parents*

222 The effect of educational interventions delivered to parents on change in their AMR knowledge,
223 attitudes or beliefs alone,^{11, 12} or in combination with parents' antimicrobial stewardship behaviour
224 outcomes^{16, 19, 23, 28} was measured in six studies. The majority of these interventions were directed to
225 parents or caregivers of children under the age of 6.^{11, 12, 19, 28} In the remaining two studies,
226 intervention was delivered to households with at least one children over 5 years old²³ and parents of
227 children aged 12-13.¹⁶

228 As shown in Table 2, all studies showed a significant increase in knowledge following the
229 interventions. In addition, four of the reviewed interventions also had a positive effect on changing
230 parents' antimicrobial stewardship behaviour. Cebotarenco & Bush (2008)¹⁶ found, that parents in
231 the intervention group were 5.2 times more likely than other parents to indicate they had not taken
232 an antibiotic for colds or flu ($p < 0.001$). In Trepka *et al.* (2001)¹⁹, the proportion of parents who
233 expected an antibiotic for their child and did not receive one declined in the intervention area from
234 14% to 9%, while it increased from 7% to 10% in the control area ($p = 0.003$) and the percentage of
235 parents reporting that they brought their child to another physician because they did not receive an
236 antibiotic decreased from 5% to 2% in the intervention area and increased from 2% to 4% in the
237 control area ($p = 0.02$). Larson *et al.* (2009)²³ found that the percentage of participants reporting

238 using alcohol hand sanitizers has increased from 1.4% to 66.8% following the intervention ($p =$
239 0.001) while the percentage of those reporting that at least one member of their household had
240 been vaccinated against influenza has increased from 63.7% to 73.9% ($p = 0.001$). Stockwell *et al.*
241 (2010)²⁸ on the other hand, demonstrated that the number of parents reporting that they sought
242 antibiotics without a prescription when their child was sick has decreased from 6 to 1 ($p = 0.06$).

243 *General public*

244 The general public were the population of interest in eight of the included studies.^{13, 14, 18, 21, 24, 25, 29, 30}
245 Apart from Curry *et al.* (2006)²¹ who used printed materials in the form of posters and leaflets, all
246 studies were mass media campaigns, including four studies that measured the effects of the national
247 campaign intervention.^{18, 21, 25, 29}

248 Five studies demonstrated a significant effect on the general public's knowledge and attitudes^{14, 21,}
249 ^{24, 25, 29} (Table 2). With respect to antimicrobial stewardship behaviour amongst the public, four
250 studies report a significant effect following the intervention.^{14, 21, 25, 29}

251 Gonzales *et al.* (2008)¹⁴ found that visits to paediatricians declined in the intervention group for all
252 conditions but mostly for acute respiratory infections ($p = 0.01$). Similarly, Curry (2006)²¹
253 demonstrated a significant decrease in the numbers of respondents who reported consulting a
254 doctor for the common cold ($p = 0.026$). The results of Wutzke, (2006)²⁹ showed that significantly
255 less participants reported using antibiotics for cough, cold or flu following the intervention (7.4%) in
256 comparison to baseline data (10.8%; percentage point change = 3.4; 95% CI: 1.3–5.5). Mazinska &
257 Hryniewicz, (2010)²⁵ on the other hand, demonstrated a significant increase in the percentage of
258 respondents who have limited the use of antibiotics (from 27% at baseline to 43% post
259 intervention), have become more disciplined and cautious in their use (from 3% to 24%), and who
260 paid attention to the correct dosage (from 6% to 18%; no p values given).

261 The remaining three studies did not show a significant positive effect on outcomes of interest.^{13, 18, 30}
262 In Mainous *et al.* (2009),³⁰ intervention designed to decrease self-medication with antibiotics
263 surprisingly resulted in significantly greater percentage of the intervention Latino community group
264 using antibiotics without a prescription in comparison with the control group (OR = 1.81; 95% CI,
265 1.02-3.22). McNulty *et al.* (2010)¹⁸ on the other hand, found no positive effect on participants'
266 knowledge or antimicrobial stewardship behaviour following a national campaign, and there was a
267 significant increase in the percentage of respondents from the intervention area who reported
268 retaining leftover antibiotics (p <0.001). Formoso *et al.* (2013)¹³ reported that knowledge
269 consistency with the national campaign messages either worsened (p <0.05) or did not improve in
270 both the intervention and control groups after the intervention.

271 **Discussion**

272 ***Main findings of this study***

273 This systematic review provides an in-depth examination of the effectiveness of interventions that
274 target the public to increase their knowledge, understanding of AMR and engagement with
275 antimicrobial stewardship behaviours. We have also identified patterns between target populations
276 and relative intervention effectiveness. The findings present a complex picture reflecting the
277 heterogeneity of the studies.

278 Our analysis has shown that interventions targeting schoolchildren and parents have notable
279 potential. All interventions that targeted schoolchildren or parents showed a significant effect on the
280 outcome of interest. However, effective school-based interventions tended to only have effects of
281 increasing knowledge. In addition, these studies measured only short-term outcomes. In contrast,
282 interventions targeting parents demonstrated changes in behaviour in addition to knowledge, with
283 the follow up period ranging from 2 weeks²⁸ to 3 years.¹²

284 With regards to the interventions targeting the general public, the picture is less clear. Although the
285 majority (n = 5) of these studies demonstrated effectiveness of interventions in improving the
286 public's AMR knowledge or their antimicrobial stewardship behaviour, three studies did not, with
287 two showing a decrease in AMR knowledge¹³ and in antimicrobial stewardship behaviour.³⁰ These
288 findings highlight the need to examine differences in the content between these interventions
289 targeting the general public.

290 Patterning of the effectiveness across the type of target population also suggests that different
291 target populations should receive different interventions with different primary outcomes.
292 Nevertheless, targeting children alone is unlikely to make a major contribution to AMR because
293 attitudes and the public's antimicrobial stewardship behaviours may be passed down through
294 generations. Thus, using the power of familial social influence and parental duty where children's
295 AMR education within school is reinforced and boosted by parental interventions in the home, might
296 be a more appropriate approach for the achievement of desired cultural change. This indicates the
297 potential of a multimodal intervention or programmatic approach to AMR related interventions.

298 An ideal approach would be to address the entire population simultaneously, yet segmenting it to
299 target sub-populations. Through such segmentation, or stratification of the general public, diverse
300 tailored interventions addressing different sub-population, would be a strategic way to begin the
301 process of cultural change required to reduce the drivers of AMR.

302 The nature of the increase in knowledge that is needed can also be specified by drawing on other
303 evidence syntheses that has shown that the public's AMR knowledge and understanding of their
304 contribution to AMR is generally poor.³ Therefore, in addition to changing the public's understanding
305 of appropriate antimicrobial use, interventions should also target the public's understanding of AMR
306 to enable the public to understand their central role in tackling AMR, and the risks for the
307 intervention recipient, their loved ones and the wider population.

308 ***Findings in relation to other research***

309 In their recent paper, Wells & Piddock (2017)³¹ argued that amongst other actions, an urgent review
310 of educational campaigns is required in order to fulfil UK and European AMR action plans. Our
311 review addresses this need. Furthermore, to our knowledge this is the first systematic review that
312 provides such an in-depth examination of the effectiveness of AMR related interventions that target
313 the public specifically.

314 Previous literature focused on the level of the public's AMR knowledge and beliefs,^{3, 4}
315 communication interventions or interventions that target both, the public and healthcare
316 professionals.⁶ The latter found that multi-component interventions improve the public's knowledge
317 of appropriate antimicrobial use, specifically in relation to antibiotics and that interventions
318 including both, physician and public education appear to be effective in reducing antibiotic use.⁶
319 Similarly, Cross *et al.* (2016)³² reported that multi-modal communication interventions targeting
320 both the public and clinicians can reduce antibiotic prescribing in high-income countries. Although,
321 our review focused on the general public population specifically, the potential of multi-faceted
322 interventions was also highlighted in our work.

323 Another previous systematic review by King *et al.* (2015)⁷ reviewed the evidence of effectiveness
324 and cost-effectiveness of interventions changing the public' risk related behaviours in relation to
325 antimicrobial use. The review showed that direct contact education interventions were consistently
326 more effective than mass media interventions.⁷ This appears to explain our findings on the varying
327 effectiveness of interventions targeting the general public, as majority of these studies used mass
328 media interventions.

329 There is also a body of evidence on large-scale antibiotic campaigns that although were not eligible
330 for inclusion in our review as the participants were both members of the public and healthcare
331 professions. A literature review showed that there have been numerous multifaceted antibiotic
332 awareness campaigns launched in high-income countries;³³ however, there was substantial

333 heterogeneity in outcomes, including knowledge and awareness, use of antibiotics and antimicrobial
334 resistance, and the interventions themselves often lack a robust grounding in behavioural and
335 social science theory. The majority of campaigns included in the review targeted both the general
336 public and healthcare professionals simultaneously and they appeared to result in a reduction of
337 antibiotic use.³³ It therefore appears that targeting different populations at the same time might
338 result in desired outcomes as healthcare professional's prescribing decisions might also be
339 influenced by the patient, while patient's behaviour might be affected by the prescriber's advice.
340 One such campaign, conducted in the UK in 2014 simultaneously targeted members of the public
341 and healthcare professionals who pledged as Antibiotic Guardians, and showed an increase in AMR
342 knowledge and commitment to pledge behaviour in both surveyed sub-populations.³⁴ Another
343 antibiotic awareness campaign conducted in Hong Kong, targeted the general public, patients and
344 healthcare professionals in a segmented fashion, and resulted in a significant improvement (p
345 ≤ 0.002) in respondents' knowledge on prudent use of antibiotics following the campaign.³⁵ Yet
346 another successful large-scale antibiotic awareness campaign segmented to target the general public
347 and healthcare professionals was conducted in France.³⁶ The effectiveness of this campaign in
348 reducing the number of antibiotic prescriptions was evaluated and showed a 26.5% (95% CI 33.5–
349 19.6) decrease in the total number of antibiotic prescriptions following the campaign, with the
350 greatest decrease of 35.8% (95% CI -48.3% to -23.2%) in prescriptions issued for children and young
351 adults in the 21-25 years age group (24.1% decrease; CI not provided).³⁶ These findings further
352 emphasise the potential of programmatic approach to AMR related interventions segmented to
353 different target sub-populations, as suggested previously in our main findings section.

354 ***Strengths and limitations***

355 We have conducted a rigorous search and systematic review accompanied by a narrative synthesis.
356 Although, similar work concerning the effectiveness of interventions aimed to improve antibiotic use
357 has been conducted previously,³² our work focused on interventions targeting the general public

358 population exclusively and did not include outcomes related to healthcare professionals' AMR
359 awareness or antimicrobial stewardship, such as antibiotic prescribing. Our analysis provides a sense
360 of what is normative within this field, what has been attempted before and what could be repeated.
361 It also provides a unique and valuable contribution to the available literature. However, the study
362 also has limitations.

363 First, because the UK Antimicrobial Resistance Strategy and Action Plan was launched by the
364 Department of Health (DH) in 2000³⁷, followed by the publication of the WHO Global Strategy for
365 Containment of Antimicrobial Resistance in 2001¹, we limited our search to publications from 2000
366 onwards. This could result in omission of important, older papers. Second, the studies from low- and
367 middle-income countries were underrepresented in our review. Thus, relevance and applicability of
368 our findings to different geographical areas or resource contexts is limited. Third, the risk of bias was
369 assessed only for studies that met the EPOC study design criteria. However, a suitable, validated tool
370 for assessing the risk of bias of non-controlled before and after studies could not be identified.
371 Furthermore, using different instrument could result in ambiguities in relation to the quality of
372 stronger designs. The overall quality of the evidence was rather low. Major problems were
373 associated with randomisation in experimental designs and the evaluation of mass media and other
374 population level interventions. As these kinds of interventions aim for maximum population reach, it
375 is difficult to attain adequate controls or indeed randomise at this population level. Therefore, good
376 quality study designs are systematically less likely to be identified within this kind of population level
377 intervention literature. Notwithstanding this, there was a considerable heterogeneity in outcomes.
378 There are no standardised ways of measuring the public's AMR related knowledge or associated
379 stewardship behaviours. Furthermore, change in knowledge, awareness or beliefs, which were the
380 most common outcome measures across the included studies, might not necessary lead to desired
381 behaviour change. As a result, it is particularly challenging to build cumulative knowledge regarding
382 the effectiveness of interventions to increase the public's engagement with antimicrobial
383 stewardship. Another limitation is that given the problems with the quality of primary research, our

384 measure of relative effectiveness should be treated with caution as this was based on our relative
385 measure and are not equivalent of a strong evidence base within typical evidence based guidance.
386 Finally, we did not conduct an analysis of the cost-effectiveness of reviewed interventions; however,
387 for the majority of studies included in our review, cost effectiveness data was not reported.

388 ***Recommendations for future research***

389 Although, our work demonstrated the potential of intervention, that targets particular sub-
390 populations of the general public, taking into account the low quality of reviewed evidence, lack of
391 cost-effectiveness evaluation and underrepresentation of studies from low- and middle-income
392 countries, these findings must be treated with caution. There is a need for well-designed,
393 randomised experimental studies focusing on behavioural outcomes of the interventions.
394 Furthermore, measures of AMR knowledge and stewardship behaviours need to be standardised
395 and there is a need for improvement of the reporting standards to ensure detailed and transparent
396 reporting of intervention components. Finally, considering the underrepresentation of studies from
397 low- and middle-income countries, there is a need for the development and evaluation of similar
398 interventions within such settings.

399 **Conclusions**

400 Although some evidence on the effectiveness of interventions that target the general public to
401 engage with the problem of AMR exists, the public's understanding of AMR and their role in
402 combating this problem remains poor. Thus, there is a need for a cultural change and effective
403 engagement of the public in addition to other key stakeholders. This need could be addressed
404 through development of well-designed AMR related interventions robustly grounding within
405 behavioural and social science theory. Our work provided an in-depth examination of the
406 effectiveness of AMR related interventions targeting the members of the public specifically. We
407 suggests that future policy makers should consider multimodal segmented population level

408 intervention that tailors its core messages to children, parents and the wider general public alike,
409 particularly in high-income geographical areas. Future interventions should convey messages that
410 elicit the public's motivation to make their own efforts to address AMR as a growing problem for all
411 and a problem for the present as much as for the future.

412 **Acknowledgements**

413 **Funding**

414 This work was supported by Health Protection Scotland (CARS_2016-06-08/RIE 15-127).

415 Glasgow Caledonian University was responsible for initiating, managing and sponsoring this

416 review.

417 **Transparency declarations**

418 The Authors declare no financial interests, commercial affiliations, or conflicts of interest.

419 **References**


- 420 1. World Health Organization 2001 *WHO global strategy for containment of antimicrobial*
421 *resistance*. WHO.
422 http://apps.who.int/iris/bitstream/10665/66860/1/WHO_CDS_CSR_DRS_2001.2.pdf.
- 423 2. O'Neill J. *Tackling Drug-Resistant Infections Globally: Final Report and*
424 *Recommendations*. [https://amr-](https://amr-review.org/sites/default/files/160525_Final%20paper_with%20cover.pdf)
425 [review.org/sites/default/files/160525_Final%20paper_with%20cover.pdf](https://amr-review.org/sites/default/files/160525_Final%20paper_with%20cover.pdf)
- 426 3. Gualano MR, Gili R, Scaiola G *et al*. General population's knowledge and attitudes about
427 antibiotics: a systematic review and meta-analysis. *Pharmacoepidemiol Drug Saf* 2015; **24**: 2-
428 10.
- 429 4. McCullough A, Parekh S, Rathbone J *et al*. A systematic review of the public's
430 knowledge and beliefs about antibiotic resistance. *J Antimicrob Chemother* 2015; **71**: 27-33.
- 431 5. Pinder R, Sallis A, Berry D *et al*. *Behaviour Change and Antibiotic Prescribing in*
432 *Healthcare Settings Literature Review and Behavioural Analysis*. PHE.
433 [https://www.gov.uk/government/publications/antibiotic-prescribing-and-behaviour-change-](https://www.gov.uk/government/publications/antibiotic-prescribing-and-behaviour-change-in-healthcare-settings)
434 [in-healthcare-settings](https://www.gov.uk/government/publications/antibiotic-prescribing-and-behaviour-change-in-healthcare-settings)
- 435 6. Haynes C, McLeod C. *A review of Reviews of Educational Interventions Designed to*
436 *Change the Public's Knowledge and Behaviour in Relation to Antimicrobial Use and*
437 *Antimicrobial Resistance that Target Healthcare Professionals and Patients*.
438 [https://www.nice.org.uk/guidance/ng63/documents/antimicrobial-resistance-changing-](https://www.nice.org.uk/guidance/ng63/documents/antimicrobial-resistance-changing-riskrelated-behaviours-in-the-general-population-evidence-review-32)
439 [riskrelated-behaviours-in-the-general-population-evidence-review-32](https://www.nice.org.uk/guidance/ng63/documents/antimicrobial-resistance-changing-riskrelated-behaviours-in-the-general-population-evidence-review-32)
- 440 7. King S, Exley J, Taylor J *et al*. Antimicrobial stewardship: the effectiveness of
441 educational interventions to change risk-related behaviours in the general population: a
442 systematic review. *Rand Health Q* 2016; **5**: 2.
- 443 8. Moher D, Liberati A, Tetzlaff J *et al*. Preferred reporting items for systematic reviews
444 and meta-analyses: the PRISMA statement. *PLoS Med* 2009; **6**: e1000097.
- 445 9. Cochrane Effective Practice and Organisation of Care (EPOC). *EPOC Resources for*
446 *Review Authors*. <http://epoc.cochrane.org/resources/epoc-resources-review-authors>
- 447 10. Ryan R, Cochrane Consumers and Communication Review Group. *Cochrane Consumers*
448 *and Communication Review Group: Data Synthesis and Analysis*.
449 [http://cccr.org.cochrane.org/sites/cccr.org.cochrane.org/files/public/uploads/meta-](http://cccr.org.cochrane.org/sites/cccr.org.cochrane.org/files/public/uploads/meta-analysis_revised_december_1st_1_2016.pdf)
450 [analysis_revised_december_1st_1_2016.pdf](http://cccr.org.cochrane.org/sites/cccr.org.cochrane.org/files/public/uploads/meta-analysis_revised_december_1st_1_2016.pdf)
- 451 11. Croft DR, Knobloch MJ, Chyou P *et al*. Impact of a child care educational intervention
452 on parent knowledge about appropriate antibiotic use. *WMJ* 2007; **106**: 78-84.
- 453 12. Huang SS, Rifas-Shiman SL, Kleinman K *et al*. Parental knowledge about antibiotic use:
454 results of a cluster-randomized, multicomponent intervention. *Pediatrics* 2007; **119**: 698-706.
- 455 13. Formoso G, Paltrinieri B, Marata AM *et al*. Feasibility and effectiveness of a low cost
456 campaign on antibiotic prescribing in Italy: community level, controlled, non-randomised trial.
457 *BMJ* 2013; **347**: f5391.
- 458 14. Gonzales R, Corbett KK, Wong S *et al*. "Get smart Colorado": impact of a mass media
459 campaign to improve community antibiotic use. *Med Care* 2008; **46**: 597-605.
- 460 15. Pontes MC, Pontes NM. Debiasing effects of education about appropriate antibiotic
461 use on consumers' preferences for physicians. *Health Care Manage Rev* 2005; **30**: 9-16.
- 462 16. Cebotarenco N, Bush PJ. Reducing antibiotics for colds and flu: a student-taught
463 program. *Health Educ Res* 2008; **23**: 146-57.

- 464 17. Lecky DM, McNulty CA, Touboul P *et al.* Evaluation of e-Bug, an educational pack,
465 teaching about prudent antibiotic use and hygiene, in the Czech Republic, France and England.
466 *J Antimicrob Chemother* 2010; **65**: 2674-84.
- 467 18. McNulty CA, Nichols T, Boyle PJ *et al.* The English antibiotic awareness campaigns: did
468 they change the public's knowledge of and attitudes to antibiotic use? *J Antimicrob Chemother*
469 2010; **65**: 1526-33.
- 470 19. Trepka MJ, Belongia EA, Chyou PH *et al.* The effect of a community intervention trial
471 on parental knowledge and awareness of antibiotic resistance and appropriate antibiotic use in
472 children. *Pediatrics* 2001; **107**: e6.
- 473 20. Azevedo MM, Pinheiro C, Yaphe J *et al.* Assessing the impact of a school intervention
474 to promote students' knowledge and practices on correct antibiotic use. *Int J Environ Res*
475 *Public Health* 2013; **10**: 2920-31.
- 476 21. Curry M, Sung L, Arroll B *et al.* Public views and use of antibiotics for the common cold
477 before and after an education campaign in New Zealand. *N Z Med J* 2006; **119**: U1957.
- 478 22. Farrell D, Kostkova P, Weinberg J *et al.* Computer games to teach hygiene: an
479 evaluation of the e-Bug junior game. *J Antimicrob Chemother* 2011; **66** Suppl 5: v39-v44.
- 480 23. Larson EL, Ferng YH, McLoughlin JW *et al.* Effect of intensive education on knowledge,
481 attitudes, and practices regarding upper respiratory infections among urban Latinos. *Nurs Res*
482 2009; **58**: 150-7.
- 483 24. Madle G, Kostkova P, Mani-Saada J *et al.* Changing public attitudes to antibiotic
484 prescribing: can the internet help? *J Innov Health Inform* 2004; **12**: 19-26.
- 485 25. Mazińska B, Hryniewicz W. Kampania edukacyjna Europejski Dzień Wiedzy o
486 Antybiotykach—czy wpłynęła na zmianę postaw społeczeństwa w Polsce *Pol Merkur Lekarski*
487 2010; **29**: 296-303.
- 488 26. McNulty CA, Bowen J, Gelb D *et al.* "The Bug Investigators": Assessment of a school
489 teaching resource to improve hygiene and prudent use of antibiotics. *Health Educ* 2007; **107**:
490 10-26.
- 491 27. McNulty CA, Swan AV, Boland D. Schools' antimicrobial resistance: National Advice to
492 the Public campaign—a pilot study. *Health Educ* 2001; **101**: 235-42.
- 493 28. Stockwell MS, Catalozzi M, Meyer D *et al.* Improving care of upper respiratory
494 infections among Latino Early Head Start parents. *J Immigr Minor Health* 2010; **12**: 925-31.
- 495 29. Wutzke SE, Artist MA, Kehoe LA *et al.* Evaluation of a national programme to reduce
496 inappropriate use of antibiotics for upper respiratory tract infections: effects on consumer
497 awareness, beliefs, attitudes and behaviour in Australia. *Health Promot Int* 2007; **22**: 53-64.
- 498 30. Mainous AG, Diaz VA, Carnemolla M. A community intervention to decrease antibiotics
499 used for self-medication among Latino adults. *Ann Fam Med* 2009; **7**: 520-6.
- 500 31. Wells V, Piddock LJV. Addressing antimicrobial resistance in the UK and Europe. *Lancet*
501 *Infect Dis* 2017; **17**: 1230-1231.
- 502 32. Cross ELA, Tolfree R, Kipping R. Systematic review of public-targeted communication
503 interventions to improve antibiotic use. *J Antimicrob Chemother* 2016; **72**: 975-87.
- 504 33. Huttner B, Goossens H, Verheij T *et al.* Characteristics and outcomes of public
505 campaigns aimed at improving the use of antibiotics in outpatients in high-income countries.
506 *Lancet Infect Dis* 2010; **10**: 17-31.
- 507 34. Chaintarli K, Ingle SM, Bhattacharya A *et al.* Impact of a United Kingdom-wide
508 campaign to tackle antimicrobial resistance on self-reported knowledge and behaviour change.
509 *BMC Public Health* 2016; **16**: 393.
- 510 35. Ho ML, Cowling BJ, Seto WH *et al.* Determinants of an effective antibiotic campaign:
511 lessons from Hong Kong. *J Glob Antimicrob Resist* 2014; **2**: 334-7.

- 512 36. Sabuncu E, David J, Bernède-Bauduin C *et al.* Significant reduction of antibiotic use in
513 the community after a nationwide campaign in France, 2002–2007. *PLoS Med* 2009; **6**:
514 e1000084.
- 515 37. Department of Health. *UK Antimicrobial Resistance Strategy and Action Plan*.
516 [http://antibiotic-action.com/wp-content/uploads/2011/07/DH-UK-antimicrobial-resistance-
strategy-and-action-plan.pdf](http://antibiotic-action.com/wp-content/uploads/2011/07/DH-UK-antimicrobial-resistance-
517 strategy-and-action-plan.pdf)

518 **Figures and Tables**

519 Figure 1. Study selection flowchart



521 Table 1. Review's eligibility criteria

	Inclusion criteria	Exclusion criteria
Design	Randomised controlled trials, non-randomised trial, interrupted time series studies, controlled before and after studies, non-controlled before and after studies and cohort studies.	-
Population	Members of the public	Participants recruited from healthcare settings
Intervention	Intervention designed to increase public antimicrobial awareness and/or to improve antimicrobial stewardship (through mass media, social marketing or printed media campaigns).	-
Comparator	Time bound, geographical controls or no exposure	-
Context	Non-healthcare settings; AMR or the public's antimicrobial stewardship	-
Outcomes	All relevant short, medium or long-term outcomes related to antimicrobial resistance and/or antimicrobial stewardship behaviours (knowledge/awareness, learning, public behavioural and cognition outcomes)	Antimicrobial prescribing
Publication date	Published after January 2000	Published before January 2000

522 Table 2. Study characteristics and results of the included studies

Study	Country	Design	Sample	Nature of intervention(s)	Outcome measures	Significant results
Azevedo et al. (2013) ²⁰	Braga, Portugal	NCBA	N = 82 school children	School based presentation followed by discussion.	Knowledge & attitudes	Knowledge of the correct use of antibiotics for bacterial diseases rather than viral diseases rose from 43% to 76% in the post-test (p <0.01). Knowledge of the risk of bacterial resistance to antibiotics from their incorrect use rose from 48% to 74% in the post-test (p <0.05).
Cebotarencu & Bush, (2008) ¹⁶	Chisinau, Moldova	CBA	N = 3586 school children & N = 2716 parents	Educational intervention about the use of antibiotics delivered by student volunteers trained as peer leaders delivered to their classmates & the classmates' parents.	Beliefs & behaviour	Students in both the intervention District & the Post-intervention phase were 3.2 (CI 2.065–4.909) times more likely than other students to indicate they had not taken an antibiotic.
Croft et al. (2007) ¹¹	Wisconsin, USA	RCT	N = 300 parents.	Distribution of printed materials to parents by child care staff; slide presentation delivered to staff.	Knowledge	In parents who were college graduates, the median knowledge scores were 7.0 at intervention centres & 6.5 at control centres (p <0.01).
Curry et al. (2006) ²¹	Auckland, New Zealand	NCBA	N = 400 general public	National campaign "Wise use of antibiotics". Posters & leaflets delivered to the public attending pharmacies.	Knowledge & attitudes and behaviour	Patients who had ever been to the doctor for a common cold significantly decreased (45% vs 62%; p = 0.0006). They were significantly less likely to feel positive about antibiotics in 2003 for the treatment of a cold (16% versus 33%, p = 0.00001). The perception that antibiotics were beneficial for cold/flu symptoms significantly reduced from 1998 to 2003 (p <0.05); the perceived benefit of antibiotics for tonsillitis increased from 83% to 91% in 2003 (p = 0.014) Significantly less people reported ever attending a doctor for a cold in 2003 Vs 1998 (45% vs 62%; p = 0.0006); & the number of people who would usually see a doctor for a cold decreased from 24% to 15% (p = 0.026).

Farrell et al. (2011) ²²	Glasgow, Gloucester and London, UK	NCBA	N = 1736 children	E-Bug web game	Knowledge & attitudes	No overall change in knowledge. Significant knowledge change in 3 out of 21 questions ($p \leq 0.02$).
Formoso et al. (2013) ¹³	Emilia-Romagna, Italy	NRT	N = 1200 general public	Local mass media campaign (posters, brochures & advertisements on local media) delivered to general population to raise awareness of inappropriate use of antibiotics.	Knowledge	After the intervention, consistency with campaign messages worsened (or did not improve) similarly in both intervention and control areas, the only exception being knowledge on the presumptive antiviral activity of antibiotics, worsening in the intervention area more than control area
Gonzales et al. (2008) ¹⁴	Colorado, USA	NRT	N = 1503 general public	Mass media intervention (outdoor & radio advertisements) delivered to general public about use of antibiotics.	Behaviour	Linear regression analysis showed a significant Net differences in monthly paediatric office visit rates between mass media & comparison communities before & after the campaign ($p = 0.01$).
Huang et al. (2007) ¹²	Massachusetts, USA	RCT	N = 3142 parents	Community based educational intervention occurred through 3 successive cold & flu seasons. Printed materials: (mailed newsletters, posters, pamphlets, & fact sheets in the waiting rooms of local paediatric providers, pharmacies, & child care centres).	Knowledge & attitudes	The proportion of parents who answered 7 of 10 knowledge questions correctly increased significantly in both intervention (from 52% to 64%; $p < 0.001$) & control (from 54% to 61%; $p < 0.01$) communities. Substantial improvements in percentage correct answers were seen for items on middle ear fluid (41% in 2000; 50% in 2003, $p < 0.001$) & the general question of whether antibiotics were needed for colds & flu (66% in 2000; 77% in 2003, $p < 0.001$)
Larson et al. (2009) ²³	Upper Manhattan, USA	NCBA	N = 422 households	Targeted Latino households. Educational materials (colouring book, pamphlets) based on knowledge, attitudes, & practices regarding prevention & treatment of upper respiratory tract infections. Program was delivered during home visits every 2 months.	Knowledge & attitudes and behaviour	After the intervention, the mean composite knowledge scores at baseline & end of study were 5.19 (SD = 1.60) & 5.91 (SD = 1.71) ($p < 0.001$), respectively. With regard to reported practices, significantly more participants after the intervention reported using alcohol hand sanitizers (1.4% baseline & 66.8% post-intervention, $p = 0.001$). Significantly more also reported that one or more members in their household had received the influenza vaccination after the intervention (63.7% at baseline & 73.9% post-intervention, $p = 0.001$).
Lecky et al. (2010) ¹⁷	Gloucestershire and London, England; Nice and Bordeaux, France; Prague	CBA	N = 2724 school students	School based educational intervention (e-bug) regarding inappropriate antibiotic use delivered to classes of 9-11-year-old (junior) & 12-15 year old (senior) students in state schools Included 45 min lesson	Knowledge	Junior school: Significant change in knowledge, & significant change in retention 6 weeks post-intervention, across countries. Little significant difference in knowledge change between intervention & control, with exception of Czech Republic. Senior school: Significant improvement in knowledge 6 weeks post

	and Ostrava, Czech Republic			hand-outs, worksheets, factsheets, interactive activity 7 a follow-up plenary question &-answer session.		intervention in Czech Republic. Significant - improvement in knowledge, & knowledge retention between control & intervention in England & Czech Republic
Madle et al. (2004)²⁴	London, UK	NCBA	N = 177 general public	Open access to the National electronic Library of Infection Antimicrobial Resistance website on the use of antibiotics & antibiotic resistance. The site comprises frequently asked questions & links to evidence based resources.	Knowledge & attitudes	Significant improvements in knowledge about the use of antibiotics & antibiotic resistance in 2 out of 3 statements: (1) "people cannot become resistant to antibiotics" (p <0.001, X2 = 60.357, 95% CI of change: 27.47 to 44.53); (2) "antibiotics do not cure most sore throats" (p <0.001, X2 = 19.22, 95% CI of change: 8.62 to 27.38). Significant changes in the scores assigned by users for 3 out of 4 statements designed to test users' attitudes to the information on the site (p ≤0.003). Expectations that antibiotics should be prescribed were significantly reduced after using the website (p <0.001). Non-HCWs continued to have higher expectations of antibiotics being prescribed than HCW (p = 0.0046 before and p = 0.0098 after using the website).
Mainous III et al. (2009)³⁰	South Carolina, USA	Prospective cohort study	N = 691 Self-identified Latinos	Mass media educational intervention (pamphlets, radio, newspapers) delivered to local Latino communities about use of antibiotics.	Behaviour	Numbers in the intervention group reporting that they had bought antibiotics without a prescription increased following the intervention compared to baseline (Chi sq test reported as significant but p value not given. The regression analysis showed the strongest predictor of purchase of antibiotics without a prescription in the previous 12 months was past purchase of antibiotics without a prescription outside the United States (OR = 5.72; 95% CI, 3.12-10.48). The regression analysis also showed the strongest predictor of likelihood of importing antibiotics into the United States was past purchase of antibiotics without a prescription outside the United States (OR = 3.01; 95% CI, 1.95-4.65).
Mazinska & Hryniewicz, (2010)²⁵	Poland	NCBA	N = 1000 general public	Mass media educational intervention (posters, leaflets, billboards, TV, cinemas, radio, press, magazines, thematic exhibitions, internet) implemented across the country.	Knowledge, attitudes & behaviour	Significant increase in the percentage of people who have limited the use of antibiotics 27-43%, have become more disciplined and cautious in their use from 3-24%, as well as pay attention to the correct dosage 6-18% (no p values given).
McNulty et al. (2001)²⁷	Gloucester, UK	NCBA	N = 38 year 5 school children.	School-based intervention to children aged 9-10years at a state school. Included two 90 minute interactive workshops	Knowledge	Before the workshops 23% & 26% knew antibiotics do not kill viruses but kill good bacteria, compared with 47% & 69% afterwards (p = 0.03 & 0.0001). 45% before & 73% after the

				entitled "Antibiotics and your good bugs".		workshops correctly answered all the questions ($p < 0.0001$). Children thought antibiotics helped hay fever, this improved significantly after the workshop (correct answer 28% before, 77% after ($p < 0.0001$). Overall score for 7 questions in the "where are bugs found" section was increased significantly from an average of 80.5% success to 93.2% success ($p = 0.0002$). The overall score improvement in the "How do bugs spread" section was significant ($p = 0.00001$).
McNulty et al. (2007) ²⁶	Gloucestershire, UK	NCBA	N = 198 year 5 and 6 school children	School based intervention. "Bug Investigators" pack about micro-organisms, hygiene & antibiotics The pack included 11 activity sheets, teachers' guide, poster & website.	Knowledge	Children's knowledge improved in all topic areas & was significant in 6 out of the 7 topic areas ($p < 0.005$). Improved knowledge was most significant for what antibiotics do & how to use them (percent improvement 27 (CI 22.8, 31.1) & 31 (CI 23.4, 37.7), respectively; & the value of our own good bugs (16 percent improvement).
McNulty et al. (2010) ¹⁸	England and Scotland, UK	CBA	N = 3718 general public	Mass media campaign about antibiotic use involving posters displayed in magazines & newspapers.	Knowledge & attitudes and behaviour	No positive effect of the campaigns.
Pontes & Pontes, (2005) ¹⁵	Mid-Atlantic region, USA	NRT	N = 105 university students	University based educational intervention (information booklet) to increase young adult consumers' preference for physicians who do not unnecessarily prescribe antibiotics for simple acute upper respiratory tract infections.	Attitudes	Exposure to the intervention significantly increased the mean preferred start date for antibiotics after initiation of an infection from 2.3-3.9 days ($p < 0.1$). Respondents' preferences were significantly greater for the physician who indicated he would not prescribe antibiotics in the intervention ($M = 4.84$) compared to control ($M = 3.91$, ($p < 0.01$).
Stockwell et al. (2010) ²⁸	New York City, USA	NCBA	N = 10 parents	Health literacy intervention regarding upper respiratory tract infection delivered to parents on a "Early Head start" programme. Involved 3x 1.5 hour interactive sessions & provision of kit for care of a child with such an infection.	Knowledge & attitudes and behaviour	The mean composite knowledge/attitude score increased from 4.1 (total possible: 10) to 6.6 ($p < 0.05$). Number of parents reporting that the last time their child was sick they sought antibiotics without a prescription instead of, or in addition to, seeing their health care provider has decreased from 6 to 1 ($p = 0.06$).
Trepka et al. (2001) ¹⁹	Northern Wisconsin, USA	CBA	N = 365 parents	Nurse educators delivered parent-oriented presentations in community organisations, distributed information pamphlets & displayed posters. Topics covered included antibiotic resistance & use.	Knowledge & attitudes and behaviour	From baseline to post intervention the percentage of parents with high antibiotic resistance awareness significantly increased in the intervention (change: 14.3%; 95% confidence interval [CI]: 6.6, 22.0) but not in the control group (change: 4.3%; 95% CI: -4.1, 12.7; $p = 0.015$). The proportion of parents who expected an antibiotic for their child & did not receive one, declined in the intervention

						area (14% to 9%), while it increased in the control area (7% to 10%). The difference between the 2 area changes was -8.4% (95% CI: -13.9,-2.8; p = 0.003). The percentage of parents in the intervention area who brought their child to another physician because they did not receive an antibiotic decreased (5% to 2%), while it increased in the control area (2% to 4%). The difference between the 2 area changes was -4.5% (95% CI: -8.0,-.9; p = 0.02).
Wutzke et al. (2006) ²⁹	Australia	NCBA	N = 6217 general public	National mass media intervention for consumers delivered during winter months in 2001, 2002, 2003, & 2004. About the inappropriate use of antibiotics for upper respiratory tract infection. Strategies included newsletters & brochures, mass media activity using billboards, television, radio & magazines & small grants to promote local community education.	Knowledge & attitudes and behaviour	There was a significant decline in those who believe taking antibiotics for cold & flu is appropriate, from 28.7% pre-programme in 2002 to 21.7% in 2004 (percentage point change = 7.0; 95% CI: 3.5–10.5). Significant decrease in self-reported use of antibiotics to treat cough, cold or flu, from 10.8% in 1999 down to 7.4% in 2004 (percentage point change = 3.4; 95% CI: 1.3–5.5).

523 Table 3. Risk of bias of studies meeting the EPOC criteria.

Study ID	Allocation - sequence generation	Allocation - concealment	Baseline primary outcome	Baseline secondary outcome	Baseline characteristics	Incomplete primary outcome data	Incomplete secondary outcome data	Blinding primary outcome data	Blinding secondary outcome	Contamination	Selective reporting primary outcome	Selective reporting secondary outcome	Other risks of bias
Cebotarenco & Bush	H	H	U	U	L	U	U	U	U	L	L	L	L
Croft <i>et al.</i> (2007) ¹¹	L	U	U	n/a	U	L	n/a	U	n/a	U	L	n/a	L
Formoso <i>et al.</i> (2013) ¹³	H	U	U	U	L	U	U	U	U	U	L	L	H
Gonzales <i>et al.</i> (2008) ¹⁴	H	U	U	L	U	U	U	U	L	H	L	L	H
Huang <i>et al.</i> (2007) ¹²	L	U	H	n/a	L	L	n/a	U	n/a	H	H	n/a	H
Lecky <i>et al.</i> (2010) ¹⁷	H	U	U	n/a	U	H	n/a	U	n/a	H	L	n/a	H
McNulty <i>et al.</i> (2010) ¹⁸	H	H	H	n/a	U	U	n/a	U	n/a	H	L	n/a	H
Pontes & Pontes	U	U	U	U	U	U	U	U	U	U	L	L	H
Trepka <i>et al.</i> (2001) ¹⁹	H	U	L	n/a	U	L	n/a	U	n/a	H	L	n/a	H

524 Table 4. Patterning of the effectiveness across the type of target population

Study	Interventions indicative of clear positive behaviour change in the desired direction	Interventions indicative of some positive behaviour change in the desired direction	Interventions indicative of positive effect on the antecedent of behavior in the desired direction	Interventions indicative of no effect on behaviour or antecedents of behaviour	Interventions indicative of negative effect on behaviour or antecedents of behaviour in a non-desired direction
Azevedo <i>et al.</i> (2013) ²⁰			School Children		
Cebotarenco & Bush, (2008) ¹⁶		School Children			
Croft <i>et al.</i> (2007) ¹¹			Parents; Child Care Facilities		
Curry <i>et al.</i> (2006) ²¹	General Public				
Farrell <i>et al.</i> (2011) ²²			School Children		
Formoso <i>et al.</i> (2013) ¹³					General Public
Gonzales <i>et al.</i> (2008) ¹⁴	General Public				
Huang <i>et al.</i> (2007) ¹²			Parents		
Larson <i>et al.</i> (2009) ²³	Parents				
Lecky <i>et al.</i> (2010) ¹⁷			School Children		
Madle <i>et al.</i> (2004) ²⁴			General Public		
Mainous <i>et al.</i> (2009) ³⁰					Latino Community, USA
Mazinska & Hryniewicz, (2010) ²⁵		General Public			
McNulty <i>et al.</i> (2001) ²⁷			School Children		
McNulty <i>et al.</i> (2007) ²⁶			School Children		
McNulty <i>et al.</i> (2010) ¹⁸				General Public	
Pontes <i>et al.</i> (2005) ¹⁵			Young Adults		
Stockwell <i>et al.</i> (2010) ²⁸	Latino Community Parents, USA				
Trepka <i>et al.</i> (2001) ¹⁹	Parents				
Wutzke <i>et al.</i> (2006) ²⁹	General Public				

525 **Supplementary data**

526 1. Supplementary Table S1. An example of search strategy applied for MEDLINE database

527 2. Supplementary Table S2. Data extraction tool designed for the purpose of the study

528 3. Supplementary Table S3. Excluded studies with rationale

529