

## **Emended descriptions of *Bacillus sporothermodurans* and *Bacillus oleronius* with the inclusion of dairy farm isolates of both species**

Heyndrickx, Marc; Coorevits, An; Scheldeman, Patsy; Lebbe, Liesbeth; Schumann, Peter; Rodríguez-Díaz, Marína; Forsyth, Gillian; Dinsdale, Anna; Heyrman, Jeroen; Logan, Niall A.; De Vos, Paul

*Published in:*  
International Journal of Systematic and Evolutionary Microbiology

*DOI:*  
[10.1099/ijs.0.026740-0](https://doi.org/10.1099/ijs.0.026740-0)

*Publication date:*  
2012

*Document Version*  
Author accepted manuscript

[Link to publication in ResearchOnline](#)

### *Citation for published version (Harvard):*

Heyndrickx, M, Coorevits, A, Scheldeman, P, Lebbe, L, Schumann, P, Rodríguez-Díaz, M, Forsyth, G, Dinsdale, A, Heyrman, J, Logan, NA & De Vos, P 2012, 'Emended descriptions of *Bacillus sporothermodurans* and *Bacillus oleronius* with the inclusion of dairy farm isolates of both species', *International Journal of Systematic and Evolutionary Microbiology*, vol. 62, no. 2, pp. 307-314. <https://doi.org/10.1099/ijs.0.026740-0>

### **General rights**

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

### **Take down policy**

If you believe that this document breaches copyright please view our takedown policy at <https://edshare.gcu.ac.uk/id/eprint/5179> for details of how to contact us.

1 **Emended descriptions of *Bacillus sporothermodurans* and *Bacillus oleronius***  
2 **with the inclusion of dairy farm isolates of both species**

3 Marc Heyndrickx<sup>1\*</sup>, An Coorevits<sup>3,4</sup>, Patsy Scheldeman<sup>1</sup>, Liesbeth Lebbe<sup>3</sup>, P. Schumann<sup>5</sup>,  
4 Marína Rodríguez-Díaz<sup>2</sup>, Gillian Forsyth<sup>2</sup>, Anna Dinsdale<sup>2</sup>, Jeroen Heyrman<sup>3</sup>, Niall A. Logan<sup>2</sup>  
5 and Paul De Vos<sup>3</sup>

6 <sup>1</sup>Institute for Agricultural and Fisheries Research (ILVO), Technology & Food Unit,  
7 Brusselsesteenweg 370, B-9090 Melle, Belgium

8 <sup>2</sup>Department of Biological and Biomedical Sciences, Glasgow Caledonian University,  
9 Cowcaddens Road, Glasgow G4 0BA, U.K.

10 <sup>3</sup>Vakgroep BFM WE10V, Laboratorium voor Microbiologie, Universiteit Gent, K. L.  
11 Ledeganckstraat 35, B-9000 Ghent, Belgium

12 <sup>4</sup>Laboratory of Biochemistry and Brewing, Faculty of Applied Engineering Sciences,  
13 University College Ghent, Schoonmeerstraat 52, 9000 Ghent, Belgium.

14 <sup>5</sup>DSMZ – Deutsche Sammlung von Mikro-organismen und Zellkulturen GmbH,  
15 Inhoffenstrasse 7 B, 38124 Braunschweig, Germany

16 Subject category: Taxonomic note

17 Running title: Emended descriptions of *B. sporothermodurans* and *B. oleronius*

18

19 Author for correspondence:

20 \*Prof. Dr. M. Heyndrickx, Institute for Agricultural and Fisheries Research (ILVO), Technology & Food Science  
21 Unit, Brusselsesteenweg 370, B-9090 Melle, Belgium. Tel: 32+ (0)9 272.30.44; Fax: 32+ (0)9 272.30.01 . e-mail:  
22 Marc.Heyndrickx@ilvo.vlaanderen.be  
23

24 *Bacillus sporothermodurans* is an industrially important micro-organism because of its ability to  
25 produce endospores which resist ultra high temperature (UHT) and industrial sterilization  
26 processes. It was described by Pettersson *et al.* (1996) based on seven genetically homogeneous  
27 isolates all from UHT-milk. *Bacillus oleronius*, the closest phylogenetic neighbor of *B.*  
28 *sporothermodurans*, was described by Kuhnigk *et al.* (1995), based on a single strain, isolated from  
29 the hindgut of the termite *Reticulitermes santonensis*. A polyphasic study of a heterogeneous  
30 collection of *B. sporothermodurans* and *B. oleronius* strains isolated from various sources and  
31 geographic origins led to an emended description of both species. Additional data presented are the  
32 results of fatty acids, quinones and/or cell wall analysis (polar lipids). DNA-DNA hybridizations  
33 confirmed 3 subgroups of strains obtained after SDS-PAGE analysis of cellular proteins as *B.*  
34 *sporothermodurans*. One named *B. sporothermodurans* strain (R-7489) was reclassified as a  
35 *Bacillus fordii* strain. The phenotypic profiles of both species were rather heterogeneous,  
36 sometimes different from the original descriptions and did not differ in a large number of  
37 characters, although *B. oleronius* generally gave stronger reactions in its positive tests than did *B.*  
38 *sporothermodurans*; the variable and weak reactions for both organisms with some substrates  
39 blurred the distinction between both. However, differences in polar lipid, SDS-PAGE and  
40 menaquinone profiles clearly allow distinction between the two species.

41 *Bacillus sporothermodurans* was described by Pettersson *et al.* (1996) as a species producing  
42 highly heat-resistant endospores that may survive ultra high temperature (UHT) treatment of  
43 milk. The spore resistance to UHT-conditions (typically 140°C for a few seconds) was proven  
44 both with an isolate from UHT-milk (Huemer *et al.*, 1998) as with spores naturally occurring  
45 in contaminated UHT-milk (Scheldeman *et al.*, 2006). Isolates of this species were also found  
46 in sterilized milk and milk products such as UHT-cream, chocolate milk, evaporated milk and  
47 reconstituted milk (Herman *et al.*, 2000). As a consequence, surviving spores may germinate  
48 and grow in the consumer milk or milk product causing the technological problem of non-  
49 sterility. This problem has been encountered by different dairy companies worldwide with the

50 remarkable observation that one clone (HRS-clone) was concerned in the majority of the cases;  
51 only a few German isolates were found to be different from this HRS-clone by a non-  
52 hierarchical three-dimensional scaling of molecular typing data (Guillaume-Gentil *et al.* 2002).  
53 In the same study, farm isolates of *B. sporothermodurans* isolated from raw milk, feed  
54 concentrate and silage were shown to be genetically very heterogeneous and different from the  
55 HRS-clone. These farm isolates were identified as *B. sporothermodurans* by a specific PCR  
56 and for an isolate from raw milk and from feed concentrate also by DNA-DNA hybridization  
57 (Scheldeman *et al.*, 2002). In the same study as well as in Vaerewijck *et al.* (2001), *Bacillus*  
58 *oleronius* was isolated from raw milk and feed concentrate. Hitherto, this species had only  
59 been isolated from the hindgut of a termite and described on the basis of one isolate from this  
60 source (Kuhningk *et al.*, 1995).  
61 Because of the observed genetic heterogeneity, the original description of *B. sporothermodurans*,  
62 which was based on only a few genetically homogeneous UHT isolates belonging to the HRS-clone  
63 (Pettersen *et al.*, 1996), may no longer be adequate. One example is the upper growth limit reported  
64 by Pettersen *et al.* (1996) to be 50 °C for the type strain but shown to be as high as 55 °C for farm  
65 isolates (Scheldeman *et al.*, 2002). This prompted us to prepare an emended description of this  
66 industrially important organism as well as of its phylogenetic neighbour, *B. oleronius*, with the  
67 inclusion of genetically heterogeneous farm isolates for both species.

68 The *B. sporothermodurans* and *B. oleronius* strains used are listed in Table 1. They were  
69 grown on Brain Heart Infusion (BHI) (Oxoid) supplemented with Bacteriological agar no. 1  
70 (15 g l<sup>-1</sup>) (Oxoid) and filter sterilized vitamin B<sub>12</sub> (1 mg l<sup>-1</sup>) at 37 °C for 48 h for all analyses  
71 (unless otherwise indicated). The *B. sporothermodurans* specific PCR (BSPO-PCR) described  
72 by Scheldeman *et al.* (2002) was carried out on all strains listed above as well as on some  
73 additional *Bacillus fordii* strains described by Scheldeman *et al.* (2004). For determination of  
74 the G + C content of some selected strains, approximately 1 gram of biomass was harvested  
75 from agar plates and DNA was purified as in Logan *et al.* (2000) with the modifications as

76 outlined by Heyndrickx *et al.* (2004). The G + C content of DNA was determined by HPLC  
77 using further specifications given by Logan *et al.* (2000). DNA-DNA hybridizations were  
78 performed for strains *B. sporothermodurans* LMG 17668<sup>T</sup>, LMG 19637 and R-6778, for  
79 *Bacillus shackletonii* LMG 18435<sup>T</sup> and for *Bacillus acidicola* DSM 14745<sup>T</sup> using a  
80 modification (Willems *et al.*, 2001) of the microplate method described by Ezaki *et al.* (1989)  
81 with a reassociation temperature of 37 °C. All strains (together with the additional *B. fordii*  
82 strains mentioned above), grown on supplemented BHI as outlined above, were subjected to  
83 SDS-PAGE analysis of whole cell proteins according to Pot *et al.* (1994). The SDS-PAGE  
84 data were collected and interpreted as described by Vauterin & Vauterin (1992). For gas  
85 chromatographic analysis of methylated cellular fatty acids (FAME), cells of the same strains  
86 were grown for 24h on supplemented BHI. Further analysis was performed as described by  
87 Dawyndt *et al.* (2006). All *B. sporothermodurans* and *B. oleronius* strains together with type  
88 (and reference) strains of unreactive or weakly reactive species were phenotypically  
89 characterized by the methods of Logan & Berkeley (1984): thirteen routine biochemical tests,  
90 and 48 tests for acid production from a range of carbohydrates were made using the API 20E  
91 and 50CHB kits (bioMérieux), following cultivation of *B. sporothermodurans* and *B. oleronius*  
92 strains on BHI agar supplemented with vitamin B<sub>12</sub> at 37 °C for 24 hours. Other phenotypic  
93 characters were determined as described by Heyrman *et al.* (2004), but using BHI or BHI agar  
94 supplemented with vitamin B<sub>12</sub> as the basal media. Phenotypic data for other species were  
95 obtained on Tryptic Soy Agar (TSA) (Oxoid). For peptidoglycan analysis, whole cell  
96 hydrolysates (4N HCl, 100°C, 16 hours) of *B. sporothermodurans* LMG 17668<sup>T</sup> and *B.*  
97 *oleronius* LMG 17952<sup>T</sup> were subjected to thin layer chromatography on cellulose plates using  
98 the solvent system of Rhuland *et al.* (1955). Quinones for both strains were determined as  
99 described by Groth *et al.* (1996). For polar lipid determination, cells of both type strains were  
100 grown for 24h at 37 °C in BHI broth (1liter flasks) supplemented with vitamin B<sub>12</sub> and  
101 analyses were carried out by the Identification Service of the DSMZ (Braunschweig,

102 Germany) by two-dimensional silica gel thin-layer chromatography and detection with  
103 appropriate reagents (Tindall *et al.*, 1990a & b).

104 Numerical analysis of SDS-PAGE patterns of whole cell proteins (supplementary Fig.) revealed  
105 three groups that all contain strains sharing at least 80 % similarity between their SDS-PAGE  
106 profiles. Groups 1 and 2 are subdivided at about 75 % similarity and group 3 branches at about 65  
107 %. The first group is quite heterogeneous and contains the *B. sporothermodurans* isolates. This  
108 group can be further subdivided in three SDS-PAGE subgroups (I to III). The first subgroup I  
109 contains sixteen strains, including the type strain, the UHT-milk strains and strains from various  
110 other sources (sterilized milk, raw milk, soy and feed concentrate). The second subgroup II  
111 contains three Belgian strains (R-6777, R-6778 and R-6779) from different sources. The third  
112 subgroup III contains five Belgian strains isolated from feed concentrate. The second main SDS-  
113 PAGE group contains different strains previously described as *B. fordii* (Scheldeman *et al.*,  
114 2004), as well as one strain (R-7489) labelled as *B. sporothermodurans*. The attribution of R-  
115 7489 to *B. sporothermodurans* is based on a positive reaction in the *B. sporothermodurans*  
116 specific PCR (BSPO-PCR) that was thought to be specific for *B. sporothermodurans*  
117 (Scheldeman *et al.*, 2002). However, the type strain of *B. fordii* (not described at the time) also  
118 gives a weak amplicon in the BSPO-PCR. On the basis of the performed analysis, R-7489 should  
119 be attributed to *B. fordii*. The third SDS-PAGE group in the SDS-PAGE clustering contains  
120 named *B. oleronius* strains and five strains putatively identified as *B. oleronius* by fatty acid  
121 analysis (Scheldeman *et al.*, 2004). SDS-PAGE thus confirmed the attribution of the latter strains  
122 to *B. oleronius*. The *B. oleronius* strains can be distinguished on the basis of their SDS-PAGE  
123 profile by a strong band in the region of 85 kDalton.

124  
125 *B. sporothermodurans* M215<sup>T</sup> (= LMG 17668<sup>T</sup>) (U49078) shares 98.2% 16S rRNA gene sequence  
126 similarity with *B. oleronius* DSM 9356<sup>T</sup> (= LMG 17952<sup>T</sup>) (X82492), 97.6% with *B. acidicola* 105-  
127 2<sup>T</sup> (= DSM 14745<sup>T</sup>) (AF547209) and 97.2% with *B. shackletonii* LMG 18435<sup>T</sup> (AJ250318). *B.*

128 *oleronius* DSM 9356<sup>T</sup> (= LMG 17952<sup>T</sup>) (X82942) shares 98% 16S rRNA gene sequence similarity  
129 with *B. acidicola* 105-2<sup>T</sup> (= DSM 14745<sup>T</sup>) and 96.6% with *B. shackletonii* LMG 18435<sup>T</sup>  
130 (AJ250318). The species status of both *B. sporothermodurans* and *B. oleronius*, showing >97%  
131 16S rRNA gene sequence similarity with the above mentioned closely related species, was  
132 confirmed by low DNA-DNA hybridisation values (< 70%) as previously determined by  
133 Scheldeman *et al.* (2002) and Albert *et al.* (2005), and as determined in this study (Table 2). DNA-  
134 DNA hybridisation values determined by Scheldeman *et al.* (2002) and during this study for *B.*  
135 *sporothermodurans* strains LMG 17833 and LMG 19620 (both belonging to SDS-PAGE subgroup  
136 I), R-6778 (SDS-PAGE subgroup II) and LMG 19637 (SDS-PAGE subgroup III) with the *B.*  
137 *sporothermodurans* type strain LMG 17668<sup>T</sup> confirmed the allocation of all three observed SDS-  
138 PAGE subgroups to the species *B. sporothermodurans* (hybridisation values > 70% (Table 2).

139  
140 Fatty acid analysis did not allow a clear distinction between *B. sporothermodurans* and *B. oleronius*  
141 (Table 3). The *B. sporothermodurans* strains showed some more heterogeneity than the *B.*  
142 *oleronius* strains (as can be derived from the greater standard deviation values), but this  
143 heterogeneity could not be linked to the heterogeneity observed in SDS-PAGE analysis. Both  
144 species could clearly be differentiated from *B. fordii* by lower amounts of the dominant fatty acid  
145 iso-C<sub>15:0</sub>, and higher amounts of anteiso-C<sub>17:0</sub>.

146 *B. sporothermodurans* strains often grew poorly and benefited from the addition of vitamin B<sub>12</sub> to  
147 the medium. They also showed weak reactions in the phenotypic tests, and these reactions were not  
148 appreciably enhanced by the addition of vitamin B<sub>12</sub> to the test media. The characters were not only  
149 weak but also variable, so that the phenotypic data do not give a clearly distinctive profile that is  
150 characteristic for the species. *B. sporothermodurans* UHT milk isolates not only grew poorly but  
151 also sporulated poorly, yet their spores showed very high heat resistances (Scheldeman *et al.*,  
152 2006). This property of resistance to high temperatures decreased with subculture in the laboratory.  
153 Sporulation was enhanced by the addition of soil extract to the BHI medium supplemented with

154  $\text{MnSO}_4$  and vitamin  $\text{B}_{12}$ , but it is not known whether such spores have enhanced heat resistance.  
155 Isolates from farm environments grew more readily than UHT milk isolates and their spores did not  
156 show very high heat resistances (Scheldeman *et al.*, 2006). Strains of *B. oleronius* did not require  
157 the growth medium to be supplemented with vitamin  $\text{B}_{12}$ , and all strains sporulated well. The  
158 phenotypic profiles of the two species did not differ in a large number of characters, although *B.*  
159 *oleronius* generally gave stronger reactions in its positive tests than did *B. sporothermodurans*; the  
160 variable and weak reactions for both organisms with some other substrates blurred the distinction  
161 between the two species (Table 4). The differentiation table includes the phylogenetic closest  
162 relatives of both species, namely *B. acidicola* and *B. shackletonii*, as well as phenotypically related  
163 species.

164 The type strains of *B. sporothermodurans* and *B. oleronius* contained both *meso*-diaminopimelic  
165 acid as diagnostic diamino acid of the cell wall peptidoglycan. This means that both species show  
166 the peptidoglycan type A1 $\gamma$  (this can be concluded from the fact that *meso*-diaminopimelic acid has  
167 been reported up to now only for the peptidoglycan type A1 $\gamma$  and for three variations of the type A4  
168  $\gamma$ ; these variations of the type A4  $\gamma$  have been found so far exclusively in the genera  
169 *Brachybacterium* and *Dermabacter*, with which *B. sporothermodurans* and *B. oleronius* have no  
170 relationship). The major polar lipids of the type strains of *B. sporothermodurans* and *B. oleronius*,  
171 LMG 17668<sup>T</sup> and LMG 17952<sup>T</sup> were diphosphatidylglycerol (DPG), phosphatidylglycerol (PG)  
172 and phosphatidylethanolamine (PE), typical for the genus *Bacillus*. Minor amounts of an  
173 aminophospholipid (PN) and an unidentified phospholipid could also be detected. However, both  
174 species can be differentiated from each other based on the presence of three other unidentified  
175 phospholipids (PL's) in *B. oleronius*.

176 The type strain of *B. sporothermodurans* contained the menaquinones MK-7, MK-8 and MK-6 (in  
177 the ratio 51 : 45 : 1), while the type strain of *B. oleronius* contained MK-7, MK-8 and MK-9 (ratio  
178 52.6 : 46.9 : 0.4).



179 Overall, it has been shown here that strains of *B. sporothermodurans* and *B. oleronius* are rather  
180 heterogeneous in phenotypic characteristics but belong to these respective valid species. Both  
181 species are difficult to differentiate from each other on the basis of biochemical and physiological  
182 characteristics. The identification of members of these species requires the use of a polyphasic  
183 approach based on well defined phenotypic, chemotaxonomic and/or genetic taxonomic properties  
184 as described above. An emended description of both species is warranted as given below, including  
185 new data on fatty acid analysis, diagnostic diamino acid, peptidoglycan type, polar lipids and  
186 menaquinone profile for *B. sporothermodurans*, and polar lipids and menaquinone profile for *B.*  
187 *oleronius*. Furthermore, some additions and modifications were made on cell morphology and spore  
188 properties and on biochemical characteristics based on API 50CHB and API 20E test results as  
189 contradictions were observed with the original species descriptions. Specifically for *B.*  
190 *sporothermodurans*, more of these tests (e.g. API 50 CHB) appeared positive in our hands.

191

192 **Emended description of *Bacillus sporothermodurans*.**

193 *Bacillus sporothermodurans* (spo.ro.ther.mo.du'rans, Gr. n. *spora*, seed, and in biology a spore;  
194 Gr. adj. *thermos*, warm, hot; L. part. adj. *durans*, resisting. N. L. part adj. *sporothermodurans*,  
195 with heat-resisting spores).

196 Aerobic, Gram-positive cells that usually occur as motile, thin rods in chains. Strains require  
197 vitamin B<sub>12</sub> for satisfactory growth. After 2d on Brain Heart Infusion (BHI) agar supplemented with  
198 5 mg l<sup>-1</sup> MnSO<sub>4</sub> and with 1 mg l<sup>-1</sup> vitamin B<sub>12</sub>, colonies are 1-2mm diameter, flat, circular, entire,  
199 beige or cream and smooth or glossy in appearance. They bear spherical to ellipsoidal endospores  
200 which lie in paracentral and subterminal, sometimes terminal, positions within slightly swollen and  
201 unswollen sporangia. Sporulation is infrequent but can be enhanced by using BHI-soil extract agar  
202 supplemented with vitamin B<sub>12</sub> and MnSO<sub>4</sub>. Spores of strains isolated from UHT milk grow poorly  
203 and sporulate poorly, but their spores show very high heat resistance and have the ability to survive

204 ultra high temperature treatment (UHT). This very high heat resistance may decrease upon  
205 subculture. Isolates from farm environments may grow more readily than UHT milk isolates but be  
206 less heat resistant. Oxidase and catalase positive. Casein and starch are not hydrolysed. In the API  
207 20E strip: nitrate is reduced to nitrite, the Voges-Proskauer reaction is variable, citrate utilisation is  
208 variable, hydrogen sulphide and indole are not produced, and the ONPG reaction is negative.  
209 Gelatin and aesculin are hydrolysed, urea is not. Growth may occur between 20 and 55 °C, with an  
210 optimum of about 37 °C. Growth occurs between pH 5 and 9, and NaCl is tolerated up to 5% (w/v).  
211 In the API 50CHB gallery, acid without gas is produced from *N*-acetyl-glucosamine, D-glucose, D-  
212 fructose, maltose, and from sucrose and D-trehalose by most strains, but reactions may be weak.  
213 Acid production from the following carbohydrates is variable: amygdalin, arbutin, D-cellobiose,  
214 gentiobiose, glycerol, mannitol, D-mannose, D-melezitose, methyl-D-glucoside, salicin, starch  
215 (weak), D-tagatose, D-turanose and xylitol (weak). Acid is not produced from the following  
216 carbohydrates: adonitol, D-arabinose, L-arabinose, D-arabitol, L-arabitol, dulcitol, erythritol, D-  
217 fucose, L-fucose, galactose, gentiobiose, gluconate, 2-keto-D-gluconate, 5-keto-D-gluconate,  
218 glycogen, inulin, lactose, D-lyxose, D-melibiose, *meso*-inositol, methyl-D-mannoside, methyl-  
219 xyloside, D-raffinose, rhamnose, ribose, sorbitol, L-sorbose and D-xylose.

220 The major cellular fatty acids (mean percentage  $\pm$  standard deviation of total fatty acids) after 24 h  
221 growth on BHI supplemented with vitamin B<sub>12</sub> at 37 °C are: iso-C<sub>15:0</sub> (33.28  $\pm$  4.42), anteiso-C<sub>15:0</sub>  
222 (24.87  $\pm$  2.32) and anteiso-C<sub>17:0</sub> (19.58  $\pm$  3.45). The following fatty acids are present in smaller  
223 amounts (mean percentage  $\pm$  standard deviation of total fatty acids): C<sub>16:0</sub> (6.69  $\pm$  2.32), iso-C<sub>16:0</sub>  
224 (6.40  $\pm$  1.61), iso-C<sub>17:0</sub> (5.29  $\pm$  1.22), iso-C<sub>14:0</sub> (1.77  $\pm$  0.70) and C<sub>14:0</sub> (1.51  $\pm$  0.81). The type strain  
225 has *meso*-diaminopimelic acid as the cell wall diagnostic diamino acid and contains the polar lipids  
226 diphosphatidylglycerol, phosphatidylglycerol, phosphatidylethanolamine, aminophospholipid and a  
227 phospholipid. Major menaquinones are MK7 and MK8. The G+C content is 36.7 mol% for the type  
228 strain LMG 17668<sup>T</sup> (=LMG 17894<sup>T</sup> (contains colony types t<sub>1</sub> and t<sub>2</sub>), = DSM 10599<sup>T</sup>, = M215<sup>T</sup>, =  
229 MB 581<sup>T</sup>) and 36.2 mol% and 36.5 mol% for strains LMG 19620 and LMG 17883, respectively;

230 the 16S rRNA gene sequence of the type strain is deposited at EMBL/GenBank under accession  
231 numbers U49078 (type I), U49079 (type II) and U49080 (type III). In the variable characters listed  
232 above, the type strain gives the following reactions: the Voges-Proskauer reaction is positive,  
233 citrate utilisation is negative, and acid without gas is produced from arbutin, D-cellobiose, glycerol,  
234 mannitol, D-melezitose, salicin, D-tagatose, D-turanose and xylitol, but not from amygdalin,  
235 gentiobiose, D-mannose, methyl-D-glucoside and starch. Spores, though scanty, are ellipsoidal,  
236 terminal, and do not swell the sporangia.

237 **Emended description of *Bacillus oleronius*.**

238 *Bacillus oleronius* (o.le.ro'ni.us, N. L. masc. adj. *oleronius*, pertaining to the island île d'Oléron  
239 (France), the place where the organism was first isolated from the hindgut of a termite).

240 Cells are non-motile, Gram-negative, medium-sized rods, that occur singly and in pairs, and  
241 sometimes form short chains of 3-4 cells. They bear ellipsoidal endospores that lie in subterminal  
242 and paracentral positions within swollen sporangia. Spores do not show very high heat resistance.  
243 After 2 d on TSA colonies are approximately 1-2 mm diameter, circular, entire, shiny, beige or  
244 cream and butyrous with slightly translucent edges. Organisms are strictly aerobic and catalase  
245 positive. Growth may occur between 30 and 50 °C, with an optimum of 37 °C. Growth occurs at pH  
246 5.7 and at 6.8. NaCl is tolerated up to 7% (w/v). Casein is not hydrolysed and starch is sometimes  
247 hydrolysed weakly. In the API 20E strip: nitrate is reduced to nitrite, the Voges-Proskauer reaction  
248 is variable, citrate is not utilized, hydrogen sulphide and indole are not produced, and the ONPG  
249 reaction is negative. Aesculin is hydrolysed, gelatin is weakly hydrolysed and urea is not  
250 hydrolysed. In the API 50CHB gallery: acid without gas is produced from *N*-acetyl-glucosamine,  
251 D-cellobiose, D-fructose, D-glucose, mannitol and D-tagatose. Acid production from the following  
252 carbohydrates is variable, and when positive it is weak: galactose, glycerol, maltose, D-mannose,  
253 ribose, salicin, starch and D-trehalose. Acid is not produced from the following carbohydrates:  
254 adonitol, amygdalin, D- and L-arabinose, D- and L-arabitol, arbutin, dulcitol, erythritol, D- and L-

255 fucose, gentiobiose, gluconate, 2-keto-D-gluconate, 5-keto-D-gluconate, methyl-D-glucoside,  
256 glycogen, *meso*-inositol, inulin, lactose, D- and L-lyxose, methyl-D-mannoside, D-melibiose, D-  
257 melezitose, rhamnose, D-raffinose, sorbitol, L-sorbose, sucrose, D-turanose, xylitol, D- and L-  
258 xylose and methyl-xyloside.

259 The major cellular fatty acids (mean percentage  $\pm$  standard deviation of total fatty acids) after 24 h  
260 growth on BHI supplemented with vitamin B<sub>12</sub> at 37 °C are: iso-C<sub>15:0</sub> (39.24  $\pm$  1.38), anteiso-C<sub>15:0</sub>  
261 (22.89  $\pm$  2.22) and anteiso-C<sub>17:0</sub> (20.78  $\pm$  0.85). The following fatty acids are present in smaller  
262 amounts (mean percentage  $\pm$  standard deviation of total fatty acids): iso-C<sub>17:0</sub> (6.76  $\pm$  1.25), iso-  
263 C<sub>16:0</sub> (5.37  $\pm$  0.50), C<sub>16:0</sub> (3.35  $\pm$  0.29) and iso-C<sub>14:0</sub> (1.09  $\pm$  0.15). The type strain has *meso*-  
264 diaminopimelic acid as the cell wall diagnostic diamino acid and contains the polar lipids  
265 diphosphatidylglycerol, phosphatidylglycerol, phosphatidylethanolamine, aminophospholipid and  
266 four different phospholipids. Major menaquinones are MK7 and MK8. The G+C content is 35.2  
267 mol% for the type strain LMG 17952<sup>T</sup> (= DSM 9356<sup>T</sup>, Rt 10<sup>T</sup>) and 34.7 mol% for LMG 17886; the  
268 16S rRNA gene sequence of the type strain is deposited at EMBL/GenBank under accession  
269 number X82492. In the variable characters listed above the type strain is positive for the Voges-  
270 Proskauer reaction, and acid without gas is produced from: glycerol, maltose, ribose, starch (weak)  
271 and D-trehalose, but not from galactose, D-mannose or salicin.

## 272 **ACKNOWLEDGEMENTS**

273 We are very grateful to Elly Engels for excellent technical assistance. We are most grateful to  
274 bioMérieux sa for providing API materials and for supporting G. F., M. R.-D. and A.D.; P. D.  
275 V. is indebted to the Fund for Scientific Research - Flanders for personnel and research grant  
276 G.0156.02. J. H. is most grateful to FWO-Vlaanderen for a personal grant. We thank J. P.  
277 Euzéby for revision of the etymology.

278

279 **REFERENCES**

- 280 **Albert, R. A., Archambault, J., Rossello-Mora, R., Tindall, B. J. & Matheny, M. (2005)**  
281 *Bacillus acidicola* sp. nov., a novel mesophilic, acidophilic species isolated from acidid  
282 Sphagnum peat bogs in Wisconsin. *Int J Syst Evol Microbiol* **55**, 2125-2130.
- 283 **Dawyndt, P., Vancanneyt, M., Snauwaert, C., De Baets, B., De Meyer, H. & Swings, J. (2006)**  
284 Mining fatty acid databases for detection of novel compounds in aerobic bacteria. *J Microbiol Meth*  
285 **66**, 410-433.
- 286 **Ezaki, T., Hashimoto, Y. & Yabuuchi, E. (1989)** Fluorometric dexoxyribonucleic acid  
287 deoxyribonucleid acid hybridization in microdilution wells as an alternative to membrane-filter  
288 hybridization in which radioisotopes are used to determine genetic relatedness among bacterial  
289 strains. *Int J Syst Bacteriol* **39**, 224-229.
- 290 **Groth, I., Schumann, P., Weiss, N., Martin, K. & Rainey, F. A. (1996)** *Agrococcus jenensis* gen.  
291 nov., sp. nov., a new genus of actinomycetes with diaminobutyric acid in the cell wall. *Int J Syst*  
292 *Bacteriol* **46**, 234-239.
- 293 **Guillaume-Gentil, O., Scheldeman, P., Marugg, J., Herman, L., Joosten, H. & Heyndrickx,**  
294 **M. (2002)** Genetic heterogeneity in *Bacillus sporothermodurans* as demonstrated by ribotyping and  
295 repetitive extragenic palindromic-PCR fingerprinting. *Appl Environ Microbiol* **68**, 4216-4224.
- 296 **Herman, L., Heyndrickx, M., Vaerewijck, M. & Klijn, N. (2000)** *Bacillus sporothermodurans* –  
297 a *Bacillus* forming highly heat-resistant spores. 3. Isolation and methods of detection. *Bull Int*  
298 *Dairy Fed* **357**, 9-14.
- 299 **Heyndrickx, M., Logan, N. A., Lebbe, L., Rodríguez-Díaz, M., Forsyth, G., Goris, J.,**  
300 **Scheldeman, P. & De Vos, P. (2004)** *Bacillus galactosidilyticus* sp. nov., an alkali-tolerant,  $\beta$ -  
301 galactosidase-producer. *Int J Syst Evol Microbiol* **54**, 617-621.

302 **Heyrman, J., Vanparrys, B., Logan, N. A., Balcaen, A., Rodríguez-Díaz, M., Felske A. & De**  
303 **Vos, P. (2004)** *Bacillus novalis* sp. nov., *Bacillus vireti* sp. nov., *Bacillus soli* sp. nov., *Bacillus*  
304 *bataviensis* sp. nov. and *Bacillus drentensis* sp. nov., five new species isolated from the Drentse A  
305 grasslands. *Int J Syst Evol Microbiol* **54**, p. 47-57

306 **Huemer, I.A., Klijn, N., Vogelsang H. W. J. & Langeveld, L. P. M. (1998)** Thermal death  
307 kinetics of spores of *Bacillus sporothermodurans* Isolated from UHT milk. *Int Dairy J* **8**, 851-855.

308 **Kuhnigk, T., Borst, E.-M., Breunig, A., König, H., Collins, M. D., Hutson, R. A. & Kämpfer,**  
309 **P. (1995)** *Bacillus oleronius* sp. nov., a member of the hindgut flora of the termite *Reticulitermes*  
310 *santonensis* (Feytaud). *Can J Microbiol* **41**, 699-706.

311 **Logan, N. A. and Berkeley, R. C. W. (1984)** Identification of *Bacillus* strains using the API  
312 system. *J Gen Microbiol* **130**, 1871-1882.

313 **Logan, N. A., Lebbe, L. Hoste, B. Goris, J. Forsyth, G. Heyndrickx, M. & De Vos P.**  
314 **(2000)** Aerobic endospore-forming bacteria from geothermal environments in northern  
315 Victoria Land, Antarctica, and Candlemas Island, South Sandwich archipelago, with the  
316 proposal of *Bacillus fumarioli* sp. nov. *Int J Syst Evol Microbiol* **50**, 1741-1753.

317 **Pettersson, B., Lembke, F., Hammer, F., Stackebrandt, E. & Priest, F. G. (1996)** *Bacillus*  
318 *sporothermodurans*, a new species producing highly heat-resistant endospores. *Int J Syst Evol*  
319 *Microbiol* **46**, 759-764.

320 **Pot, B., Vandamme, P. & Kersters, K. (1994)** Analysis of electrophoretic whole organism  
321 protein fingerprints. In *Chemical Methods in Prokaryotic Systematics*, pp. 493-521. Edited by  
322 M. Goodfellow & A. G. O'Donnell. Chichester: Wiley.

323 **Rhuland, L.E., Work, E., Denman, R.F. & Hoare, D.S. (1955)**. The behavior of the isomers  
324 of  $\alpha$ ,  $\epsilon$  diaminopimelic acid on paper chromatograms. *J Am Chem Soc* **77**, 4844-4846.

325 **Scheldeman, P., Herman, L., Goris, J., De Vos, P. & Heyndrickx, M. (2002)** Polymerase  
326 chain reaction identification of *Bacillus sporothermodurans* from dairy sources. *J Appl*  
327 *Microbiol* **92**, 983-991.

328 **Scheldeman, P., Rodríguez-Díaz, M., Goris, J., Pil, A., De Clerck, E., Herman, L., De Vos, P.,**  
329 **Logan, N. A. & Heyndrickx, M. (2004)** *Bacillus farraginis* sp. nov., *Bacillus fortis* sp. nov., and  
330 *Bacillus fordii* sp. nov., isolated at dairy farms. *Int J Syst Evol Microbiol* **54**, 1355-1364.

331 **Scheldeman, P., Herman, L., Foster, S. & Heyndrickx, M. (2006)** *Bacillus*  
332 *sporothermodurans* and other highly heat-resistant spore formers in milk. *J Appl Microbiol*  
333 **101**, 542-555.

334 **Tindall, B.J. (1990a)**. Lipid composition of *Halobacterium lacusprofundi*. *FEMS Microbiol Lett*  
335 **66**, 199-202.

336 **Tindall, B.J. (1990b)**. A comparative study of the lipid composition of *Halobacterium*  
337 *saccharovororum* from various sources. *Syst Appl Microbiol* **13**, 128-130.

338 **Vaerewijck, M., De Vos, P., Lebbe, L., Scheldeman, P., Hoste, B. & Heyndrickx, M. (2001)**  
339 Occurrence of *Bacillus sporothermodurans* and other aerobic spore-forming species in feed  
340 concentrate for dairy cattle. *J Appl Microbiol* **91**, 1074-1084.

341 **Vauterin, L. & Vauterin, P. (1992)** Computer aided objective comparison of electrophoretic  
342 patterns for grouping and identification of microorganisms. *Eur Microbiol* **1**, 37-41.

343 **Willems, A., Doignon-Bourcier, F., Goris, J., Coopman, R., de Lajudie, P., De Vos, P. &**  
344 **Gillis, M. (2001)** DNA-DNA and hybridization study of *Bradyrhizobium* strains. *Int J Syst*  
345 *Evol Microbiol* **51**, 1315-1322.

346

347 Table 1: *B. sporothermodurans* and *B. oleronius* strains used and their origin.

348	Original species	Eventual	LMG	R-number <sup>*</sup>	Other designations <sup>†</sup>	Source
349	identification	new species	number			
350		identification				
351						
352	<i>B. sporothermodurans</i>		17668 <sup>T</sup>	1948	LMG 17894 <sup>T</sup> (t <sub>1</sub> and t <sub>2</sub> ) <sup>‡</sup>	UHT-milk, Italy
353					M215 <sup>T</sup> , MB 581 <sup>T</sup> ,	
354					DSM 10599 <sup>T</sup>	
355	<i>B. sporothermodurans</i>		17897	1951	MB 372	UHT-milk, Germany
356	<i>B. sporothermodurans</i>		18461		MB 512	UHT-milk, Belgium
357	<i>B. sporothermodurans</i>		18465		MB 373	UHT-milk, Germany
358	<i>B. sporothermodurans</i>			17269	MB 1320	sterilized milk, Dominican republic
359	<i>B. sporothermodurans</i>		17883	1937	MB 385	raw milk, Belgium
360	<i>B. sporothermodurans</i>		19637	3247	MB 1668	feed concentrate, Belgium
361	<i>B. sporothermodurans</i>			6645	MB 1493	feed concentrate, Belgium
362	<i>B. sporothermodurans</i>			6662	MB 1494	soy, Belgium
363	<i>B. sporothermodurans</i>			6665	MB 1495	soy, Belgium
364	<i>B. sporothermodurans</i>			6689	MB 1496	feed concentrate, Belgium
365	<i>B. sporothermodurans</i>			6701	MB 1497	feed concentrate, Belgium
366	<i>B. sporothermodurans</i>			6710	MB 1498	feed concentrate, Belgium
367	<i>B. sporothermodurans</i>			6713	MB 1499	feed concentrate, Belgium
368	<i>B. sporothermodurans</i>			6723	MB 1500	feed concentrate, Belgium
369	<i>B. sporothermodurans</i>			6725	MB 1501	feed concentrate, Belgium
370	<i>B. sporothermodurans</i>			6744	MB 1502	feed concentrate, Belgium
371	<i>B. sporothermodurans</i>			6777	MB 1503	soy, Belgium
372	<i>B. sporothermodurans</i>			6778	MB 1504	pulp, Belgium
373	<i>B. sporothermodurans</i>			6779	MB 1505	silage, Belgium
374	<i>B. sporothermodurans</i>			6786	MB 1506	feed concentrate, Belgium
375	<i>B. sporothermodurans</i>			7342	MB 1508	feed concentrate, Belgium
376	<i>B. sporothermodurans</i>	<i>B. fordii</i>		7489	MB 1509	feed concentrate, Belgium
377	<i>B. sporothermodurans</i>		19617	3227	MB 1316	feed concentrate, Belgium
378	<i>B. sporothermodurans</i>		19620	3299	MB 1317	feed concentrate, Belgium
379	<i>B. oleronius</i>		17882	1936	MB 382	raw milk, Belgium
380	<i>B. oleronius</i>		17884	1938	MB 386	raw milk, Belgium



381	<i>B. oleronius</i>		17886	1940	MB 397	raw milk, Belgium
382	<i>B. oleronius</i>		17887	1941	MB 398	raw milk, Belgium
383	<i>B. oleronius</i>		17952 <sup>T</sup>		DSM 9356 <sup>T</sup>	termite
384	<i>B. oleronius</i>		19619	3297	MB 1318	feed concentrate, Belgium
385	<i>B. oleronius</i>			6450		milk installation, Belgium
386	<i>B. oleronius</i>			6691		feed concentrate, Belgium
387	<i>B. oleronius</i>			7770	MB 2102	milk installation, Belgium
388	<i>Bacillus</i> sp. <sup>§</sup>	<i>B. oleronius</i>	6724			feed concentrate, Belgium
389	<i>Bacillus</i> sp. <sup>§</sup>	<i>B. oleronius</i>	6464			filter cloth, Belgium
390	<i>Bacillus</i> sp. <sup>§</sup>	<i>B. oleronius</i>	6955		MB 2101	milk installation, Belgium
391	<i>Bacillus</i> sp. <sup>§</sup>	<i>B. oleronius</i>	6463			filter cloth, Belgium
392	<i>Bacillus</i> sp. <sup>§</sup>	<i>B. oleronius</i>	6504			milk installation, Belgium
393	<i>B. fordii</i>		22080 <sup>T</sup>	7190 <sup>T</sup>	DSM 16014 <sup>T</sup>	raw milk, Belgium
394					MB 1507 <sup>T</sup>	

395 \*: R, Research collection of the LMG Bacteria Culture Collection; †: MB, collection of the molecular bacteriology lab of  
396 the Institute of Agricultural & Fisheries Research (ILVO), Technology and Food Science Unit, Melle, Belgium; ‡: colony  
397 types; §: strains putatively identified as *B. oleronius* by FAME (Scheldeman *et al.*, 2004).

398 Table 2: DNA-DNA hybridisation values for *Bacillus sporothermodurans* strains and closest  
 399 relatives.

			LMG 17668 <sup>T</sup>	LMG 18435 <sup>T</sup>	DSM 14745 <sup>T</sup>	LMG 17952 <sup>T</sup>
<i>Bacillus sporothermodurans</i>	SDS-PAGE subgroup I	LMG 17668 <sup>T</sup>	100%			
		LMG 17833	81%*			
		LMG 19620	88%*			
	SDS-PAGE subgroup II	R-6778	83%			
	SDS-PAGE subgroup III	LMG 19637	76%			
<i>Bacillus shackletonii</i>		LMG 18435 <sup>T</sup>	42%	100%		
<i>Bacillus acidicola</i>		DSM 14745 <sup>T</sup>	35% <sup>†</sup>	25%	100%	
<i>Bacillus oleronius</i>		LMG 17952 <sup>T</sup>	16%*	ND	32% <sup>†</sup>	100%

400 \*: data from Scheldeman *et al.* (2002); <sup>†</sup>: data from Albert *et al.* (2005)

401 **Table 3:** Cellular fatty acid methyl ester profiles of *B. sporothermodurans*, *B. oleronius* and *B.*  
 402 *fordii*. Presented values are averages for the number of strains analysed (mentioned between  
 403 brackets) with the corresponding standard deviation.

Fatty acids	<i>B. sporothermodurans</i> (n = 24)	<i>B. oleronius</i> (n = 14)	<i>B. fordii</i> (n = 2)
Saturated fatty acids			
C <sub>14:0</sub>	1.51 ± 0.81	< 1.00	< 1.00
C <sub>16:0</sub>	6.69 ± 2.32	3.35 ± 0.29	2.65 ± 0.10
Branched fatty acids			
iso-C <sub>14:0</sub>	1.77 ± 0.70	1.09 ± 0.15	< 1.00
iso-C <sub>15:0</sub>	33.28 ± 4.42	39.24 ± 1.38	47.19 ± 0.86
anteiso-C <sub>15:0</sub>	24.87 ± 2.32	22.89 ± 2.22	25.55 ± 5.49
iso-C <sub>16:0</sub>	6.40 ± 1.61	5.37 ± 0.50	2.10 ± 1.44
iso-C <sub>17:0</sub>	5.29 ± 1.22	6.76 ± 1.25	7.65 ± 4.19
anteiso-C <sub>17:0</sub>	19.58 ± 3.45	20.78 ± 0.85	12.65 ± 2.12

404  
405

406

407 **Table 4.** Some characters for distinguishing between *B. sporothermodurans*, *B. oleronius*, *B. fordii*,  
 408 *B. acidicola*, *B. shackletonii* and some other weakly reacting species. All data were obtained in the  
 409 course of this study or have been taken from other studies in the authors' laboratories. All characters  
 410 were determined using tests in API 20E and API50CHB systems with the exception of microscope  
 411 observations, starch and casein hydrolysis.  
 412  
 413 +, > 85% positive; (+), 75-84% positive; v, variable (26-74% positive); -, 0-15% positive; w, weak  
 414 positive reaction; +/w, positive or weak positive reaction; v/w, variable and when positive the  
 415 reaction is weak.

	<i>B. sporothermodurans</i>	<i>B. oleronius</i>	<i>B. rufis</i>	<i>B. circulans</i> T	<i>B. lentus</i> T	<i>B. firmus</i>	<i>B. fordii</i>	<i>B. acidicola</i>
Motility	v	-	+	+	+	+	+	+
Spores observed	rarely	+	+	+	+	+	+	+
Spores spherical	v	-	-	-	-	-	-	-
Sporangia swollen	v	+	+	+	-	-	v	+
Anaerobic growth	-	-	+	+	-	+	-	-
Gram reaction	+	-	+	+	+	+	-	+
Starch hydrolysis	-	v/w	+	+	+	+	-	-
Casein hydrolysis	-	-	-	w	-	+	-	-
<b>API 20E tests</b>								
ONPG reaction	-	-	+	+	+	-	-	-
Citrate utilisation (Simmons')	v	-	-	-	-	-	-	+
Urease production	-	-	-	-	+	-	-	-
Voges-Proskauer test	v	v/w	-	w	-	-	-	-
Gelatin hydrolysis	+	+/w	-	-	-	+	-	-
Nitrate reduction	+	+	+	+	+	v	-	+
<b>API 50 CHB tests</b>								
Glycerol	v	v/w	-	+	-	+	-	-
L-arabinose	-	-	+	+	w	-	-	-
Ribose	-	v/w	+	w	w	w	-	+
D-xylose	-	-	+	+	w	v	-	+
Methyl-xyloside	-	-	-	+	-	-	-	-
Galactose	-	v/w	v	+	w	-	-	+
D-mannose	v	v/w	+	+	+	-	-	w
Meso-inositol	-	-	-	+	w	-	-	-

mannitol	v	+	v	+	-	+	-	w
sorbitol	-	-	-	+	w	-	-	-
Methyl-D-glucoside	v	-	v	+	+	-	-	+
Amygdalin	v	-	-	+	w	-	-	+
Arbutin	v	-	-	+	w	-	-	+
Salicin	v	v/w	v	+	w	-	-	+
D-cellobiose	v	+	v	+	w	-	-	+
Maltose	+	v/w	v	+	w	+	-	+
Lactose	-	-	+	+	+	-	-	w
D-melibiose	-	-	+	+	w	-	-	+
Sucrose	(+)	-	+	+	+	+	-	+
D-trehalose	(+)	v/w	+	+	+	+	-	+
Inulin	-	-	v	+	-	-	-	+
D-melezitose	v	-	+	+	w	-	-	w
D-raffinose	-	-	v	+	w	-	-	+
Glycogen	-	-	v	+	-	+	-	+
Xylitol	v/w	-	-	+	-	-	-	w
Gentiobiose	v	-	-	+	w	-	-	+
D-turanose	v	-	-	+	w	-	-	-

416

417

418 **Legend Supplementary figure.** Normalized computer profiles from SDS-PAGE analyses of whole  
419 cell proteins of isolates belonging to the species *Bacillus sporothermodurans*, *B. oleronius* and *B.*  
420 *fordii*. The dendrogram is based on UPGMA clustering of the correlation coefficient ( $r$ ) of the total  
421 protein profiles. The zone used for clustering is indicated in gray on the kDalton bar above the  
422 profiles.  
423  
424

