

How should we deliver sexual health services in the 2020s?

Ong, Jason J.; Estcourt, Claudia; Tucker, Joseph D.; Golden, Matthew R.; Hocking, Jane S.; Fairley, Christopher K.

Published in:
Sexual Health

DOI:
[10.1071/SH22107](https://doi.org/10.1071/SH22107)

Publication date:
2022

Document Version
Publisher's PDF, also known as Version of record

[Link to publication in ResearchOnline](#)

Citation for published version (Harvard):

Ong, JJ, Estcourt, C, Tucker, JD, Golden, MR, Hocking, JS & Fairley, CK 2022, 'How should we deliver sexual health services in the 2020s?', *Sexual Health*, vol. 19, no. 4, pp. 233-235. <https://doi.org/10.1071/SH22107>

General rights

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

Take down policy

If you believe that this document breaches copyright please view our takedown policy at <https://edshare.gcu.ac.uk/id/eprint/5179> for details of how to contact us.

How should we deliver sexual health services in the 2020s?

Jason J. Ong^{A,B,C,*} , Claudia Estcourt^D, Joseph D. Tucker^{C,E} , Matthew R. Golden^F, Jane S. Hocking^G  and Christopher K. Fairley^{A,B} 

For full list of author affiliations and declarations see end of paper

***Correspondence to:**

Jason J. Ong
Melbourne Sexual Health Centre,
580 Swanston Street, Carlton, Vic. 3053
Australia
Email: Jason.Ong@monash.edu

Handling Editor:

Roy Chan

Received: 23 June 2022
Accepted: 30 June 2022
Published: 29 August 2022

Cite this:

Ong JJ *et al.* (2022)
Sexual Health, **19**(4), 233–235.
doi:[10.1071/SH22107](https://doi.org/10.1071/SH22107)

© 2022 The Author(s) (or their employer(s)). Published by CSIRO Publishing.

This is an open access article distributed under the Creative Commons Attribution-NonCommercial-NoDerivatives 4.0 International License (CC BY-NC-ND).

OPEN ACCESS

This Special Issue of *Sexual Health* compiles the latest evidence regarding how we can optimise sexual health services in the 2020s. Despite multiple attempts to control sexually transmitted infections (STI), globally, there are still an estimated one million newly diagnosed bacterial STI each day,¹ leading to significant health, social and economic impact. If we are to reverse the course of rising trends of STIs, we must consider the broader perspective and innovate. A fundamental shift in approach is needed.

The COVID-19 pandemic has forced rapid adaptations in the delivery of health services globally – particularly in accelerating the implementation of decentralised services, patient self-care and digital health interventions. This offers us the opportunity to reconsider how sexual health service delivery should look in the 2020s; what types of services should we continue to invest in, and are there alternate models that should be explored further? Undergirding this adaptation to the new normal, one principle must remain forefront – access to comprehensive, high-quality, and appropriately targeted sexual health services is still one of the cornerstones of controlling STIs. Fairley and colleagues' review provides an overview of the effectiveness of strategies to control STIs and concludes that accessible health care is the most powerful tool to control STIs and therefore should form part of all comprehensive STI control programs.²

Do we still need specialist sexual health services?

We acknowledge that neither specialist sexual health services nor speciality pathways for workforce development are universally available. Further, in countries where specialist sexual health services exist, they may face threats of being defunded as governments push to decentralise sexual health services to primary care and non-clinical settings. Ramchadi and colleagues' review stresses the importance of sexual health services to provide care for priority populations and people who are uninsured, demonstrating that they remain the preferred service for STI care.³ In this increasingly resource-constrained environment, Medland *et al.*⁴ present a strong case for the need for sexual health services in physical locations which concentrates funding, infrastructure and expertise. These centres can also provide support to other clinical services such as general practitioners, who contribute to the provision of accessible clinical services. The authors argue that sexual health care needs are rising both in volume and complexity, not all of which can be adequately met through streamlined, decentralised, off-site care. Similarly, Woodward and colleagues⁵ retrospective audit in a community health setting in Australia explored how a physically co-located endocrine clinic (with a focus on trans and gender diverse services) with a sexual health service facilitated synergies in providing gender-affirming care and STI testing.

If we are to advocate for specialist sexual health services, what else can we do to improve their efficiency and convince funders for ongoing investment?

Getting the right people to access sexual health services

Tailoring sexual health services for the right subpopulations with the right interventions is key to using our finite resources more efficiently. Traeger and Stoové⁶ address the issue of

identifying groups most at risk for STIs to prioritise resources. The authors challenge us to rethink how we define risk as determined by demographics and sexual behaviours, demonstrating how challenging this can be because of the ongoing stigma associated with STIs, undisclosed risk behaviour, and diversification of STI epidemics beyond traditional risk groups characterised by demographics and sexual behaviours.

Once we identify key populations, how do we increase their chance of attending sexual health services? Clarke and colleagues⁷ systematic review describes the effectiveness of 13 interventions to increase attendance at pre-booked sexual health clinic appointments. They report that the most common strategies used were information about health consequences, the use of prompts/cues, and information provided by a credible source. However, these strategies were present in both effective and ineffective interventions, highlighting the need for further research to understand why these strategies work or do not work in different contexts. Tan and colleagues⁸ review gather evidence for participatory (bottoms-up) approaches to developing and implementing sexual health services. The authors discuss how community participation in clinical STI services has been operationalised and the various aspects of clinical STI services in which participatory processes have been implemented. These data have implications for enhancing community participation within clinical STI services, a neglected topic. Kularadhan *et al.*'s key informant interviews of sexual health service providers in Australia outline the strategies to improve testing among priority populations and future strategies to improve service delivery.⁹ The authors document how Australian services have expanded traditional service models and implemented new approaches to optimise service delivery.

Strengthening current services

Attendance at sexual health services provides a gateway to offer primary prevention services (to optimise sexual health) and secondary prevention services (to prevent reinfections). Jayes *et al.*¹⁰ present the strengths and weaknesses of delivering primary prevention interventions in sexual health services. This includes education and awareness building, condom promotion, and the provision of vaccines (human papillomavirus and hepatitis A and B). Golden *et al.*¹¹ argue for the need to modernise the partner notification (PN) system if we are to create meaningful change. The authors provide a valuable summary of the evidence for various approaches to PN in high-income countries, provide key principles for best practice and list priority research questions. Specifically, they show that we have failed to consistently bring innovations to scale (e.g. expedited partner therapy), define the value of new technologies in promoting PN, and modify PN to

capitalise on opportunities to achieve broader public health objectives (e.g. promotion of HIV pre-exposure prophylaxis).

Mycoplasma genitalium complex management remains a unique challenge for sexual health services globally. Sweeney and colleagues¹² review provides a synthesis of the latest evidence on the emergence of antimicrobial resistance and its practical implications on clinical management. They underscore the need for resistance-guided treatment to prevent misuse of antimicrobials. This issue highlights the broader challenges for antibiotic stewardship, related to use of empiric therapy and high frequency of STI testing.

Smarter use of technologies to enhance sexual health services

We must be constantly alert to opportunities to provide better services to our clients. The use of digital technologies to enhance the client's experience of sexual health services is an ever-developing, fascinating space. Whilst we must be wary of leaving people behind due to the digital divide (i.e. lack of meaningful access to digital technology), it is exciting to ponder how digital technologies could facilitate a fundamental shift in our approach to STI control and modernise our services to reach the right people in a faster, more accessible way.

Tucker and colleagues¹³ timely review of digital STI and HIV services uses the prevention and care continuum as a framework to demonstrate how digital services can be interwoven into existing clinical pathways or be standalone. Gibbs and colleagues¹⁴ review describe the opportunities and challenges at the individual, service and population level of measuring and evaluating sexual health using existing digital technologies. Importantly, they provide helpful recommendations informed by a social justice lens to ensure digital health benefits all. Sha and colleagues¹⁵ trial in China offers an example of how social network-based strategies can promote the uptake of HIV/syphilis self-testing kits among men who have sex with men.

Ongoing challenges for Indigenous peoples

A special mention must be made for the ongoing challenges of Indigenous populations who still carry a disproportionate STI burden compared to non-Indigenous people. Sexual health service delivery to Indigenous populations is unique where traditional barriers (including those related to structural and social determinants of health) to accessing sexual health services are compounded by additional barriers such as historical trauma. Leston and colleagues¹⁶ review presents a case study of American Indian/Alaska Native populations in the US, where the authors describe their sexual health

behaviours and review associated STI epidemiology, clinical services, and public health interventions for these populations. The authors provide a valuable list of resources and publications and examples of sexual health services (albeit inadequate) to provide culturally relevant sexual health and STI interventions. Similarly, Sianturi and colleagues¹⁷ qualitative study of indigenous Papuans in Indonesia offers critical insights into improving their HIV programs in a setting where HIV prevalence is 24 times higher than in other islands in Indonesia. The authors highlighted missing elements in current HIV programs – sensitivity towards culture-religion concepts, dealing with modernisation-integrated HIV programs and stigma reduction.

We hope our readers enjoy the compendium of papers in this Special Issue that are thought-provoking and provide practical, evidence-based, and actionable insights to optimise sexual health services in the 2020s.

Reference

- Rowley J, Vander Hoorn S, Korenromp E, Low N, Unemo M, Abu-Raddad LJ, *et al*. Chlamydia, gonorrhoea, trichomoniasis and syphilis: global prevalence and incidence estimates, 2016. *Bull World Health Organ* 2019; 97(8): 548–62P. doi:10.2471/BLT.18.228486
- Fairley CK, Chow EPF, Simms I, Hocking JS, Ong JJ. Accessible health care is critical to the effective control of sexually transmitted infections. *Sex Health* 2022; 19: 255–64. doi:10.1071/SH22024
- Ramchandani MS, Bourne C, Barbee LA, *et al*. The need for sexual health clinics, their future role, and contribution to public health. *Sex Health* 2022; 19: 346–56. doi:10.1071/SH22087
- Medland N, Taylor R, Saunders J, *et al*. Why sexual health clinics are still important in the 2020s. *Sex Health* 2022; 19: 329–35. doi:10.1071/SH22078
- Woodward S, Luu J, Mesure J, Wynne K. A collaborative model aligning adult sexual health and endocrine gender health services. *Sex Health* 2022; 19: 386–90. doi:10.1071/SH22027
- Traeger M, Stoové M. Why risk matters for STI control: who are those at greatest risk and how are they identified? *Sex Health* 2022; 19: 265–77. doi:10.1071/SH22053
- Clarke R, Heath G, Ross JDC, Farrow C. Increasing attendance at pre-booked sexual health consultations: a systematic review. *Sex Health* 2022; 19: 236–47. doi:10.1071/SH21245
- Tan RKJ, Marley G, Kpokiri EE, Wang T, Tang W, Tucker JD. Participatory approaches to delivering clinical sexually transmitted infections services: a narrative review. *Sex Health* 2022; 19: 299–308. doi:10.1071/SH22039
- Kularadhan V, Fairley CK, Chen M, *et al*. Optimising the delivery of sexual health services in Australia: a qualitative study. *Sex Health* 2022; 19: 376–85. doi:10.1071/SH22024
- Jayes D, Merrick R, Pulford C, Buitendam E, Mohammed H, Saunders J. What is the role of sexual health services in the delivery of primary prevention of sexually transmitted infections? A narrative review. *Sex Health* 2022; 19: 319–28. doi:10.1071/SH22047
- Golden MR, Gibbs J, Woodward C, Estcourt CS. Priorities in the implementation of partner services for HIV/STI in high-income nations: a narrative review of evidence and recommendations. *Sex Health* 2022; 19: 309–18. doi:10.1071/SH22060
- Sweeney EL, Whitley DM, Murray GL, Bradshaw CS. *Mycoplasma genitalium*: enhanced management using expanded resistance-guided treatment strategies. *Sex Health* 2022; 19: 248–54. doi:10.1071/SH22012
- Tucker JD, Hocking J, Oladele D, Estcourt C. Digital sexually transmitted infection and HIV services across prevention and care continuums: evidence and practical resources. *Sex Health* 2022; 19: 278–85. doi:10.1071/22023
- Gibbs J, Solomon D, Jackson L, *et al*. Measuring and evaluating sexual health in the era of digital health: challenges and opportunities. *Sex Health* 2022; 19: 336–45. doi:10.1071/SH22068
- Sha Y, Xiong Y, Ong JJ, Wang Y, Cheng M, Ni Y, Lu Y, Tucker JD, Tang W. Comparing the effectiveness of secondary distribution of HIV/syphilis dual self-testing to testing card referral in promoting HIV testing among gay, bisexual, and other men who have sex with men in Guangzhou, China: a quasi-experimental study. *Sex Health* 2022; 19: 357–66. doi:10.1071/SH21176
- Leston J, Wenger H, Reilley B, Craig Rushing S, Rink E, Warren H, Howe J, Bloomquist P, Tah T, Jeffries I, Iralu J, Thorpe P, Apostolou A, Taylor MM. Creating a path forward: understanding the context of sexual health and sexually transmitted infections in American Indian/Alaska Native populations – a review. *Sex Health* 2022; 19: 286–98. doi:10.1071/SH22040
- Sianturi EI, Latifah E, Soltief SN, Sihombing RB, Simaremare ES, Effendy C, Probandari A, Suryawati S, Taxis K. Understanding reasons for lack of acceptance of HIV programs among indigenous Papuans: a qualitative study in Indonesia. *Sex Health* 2022; 19: 367–75. doi:10.1071/SH21206

Conflicts of interest. Jason J. Ong is the Special issues Editor for *Sexual Health* and Christopher K. Fairley is the Editor-in-Chief of *Sexual Health*, both were blinded from the peer review process for this paper. All other authors declare no conflicts of interest.

Author affiliations

^AMelbourne Sexual Health Centre, Alfred Health, Melbourne, Vic., Australia.

^BCentral Clinical School, Monash University, Melbourne, Vic., Australia.

^CFaculty of Infectious Diseases, London School of Hygiene and Tropical Medicine, London, UK.

^DGlasgow Caledonian University, Glasgow, Scotland, UK.

^EUniversity of North Carolina at Chapel Hill, Chapel Hill, NC, USA.

^FDepartment of Medicine, University of Washington, Seattle, WA, USA.

^GCentre for Epidemiology and Biostatistics, Melbourne School of Population and Global Health, The University of Melbourne, Melbourne, Vic., Australia.