

Health and sociodemographic determinants of excess mortality in Spanish nursing homes during the COVID-19 pandemic: a 2-year prospective longitudinal study

Escriba-Salvans, Anna; Jerez-Roig, Javier ; Farres-Godayol, Pau; Bezerra de Souza , Dyego Leandro; Skelton, Dawn A.; Minobes-Molina, Eduard

Published in:
Zeitschrift für Gerontologie und Geriatrie

DOI:
[10.1007/s00391-024-02294-4](https://doi.org/10.1007/s00391-024-02294-4)

Publication date:
2024

Document Version
Publisher's PDF, also known as Version of record

[Link to publication in ResearchOnline](#)

Citation for published version (Harvard):
Escriba-Salvans, A, Jerez-Roig, J, Farres-Godayol, P, Bezerra de Souza , DL, Skelton, DA & Minobes-Molina, E 2024, 'Health and sociodemographic determinants of excess mortality in Spanish nursing homes during the COVID-19 pandemic: a 2-year prospective longitudinal study', *Zeitschrift für Gerontologie und Geriatrie*.
<https://doi.org/10.1007/s00391-024-02294-4>

General rights

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

Take down policy

If you believe that this document breaches copyright please view our takedown policy at <https://edshare.gcu.ac.uk/id/eprint/5179> for details of how to contact us.

Z Gerontol Geriat
<https://doi.org/10.1007/s00391-024-02294-4>
Received: 26 January 2024
Accepted: 6 March 2024

© The Author(s) 2024



Health and sociodemographic determinants of excess mortality in Spanish nursing homes during the COVID-19 pandemic: a 2-year prospective longitudinal study

Anna Escribà-Salvans^{1,2} · Javier Jerez-Roig^{1,2,3} · Pau Farrés-Godayol^{1,2} · Dyego Leandro Bezerra de Souza⁴ · Dawn A. Skelton⁵ · Eduard Minobes-Molina^{1,2}

¹ Research group on Methodology, Methods, Models and Outcomes of Health and Social Sciences (M3O). Faculty of Health Sciences and Welfare. Centre for Health and Social Care Research (CESS), University of Vic-Central University of Catalonia (UVic-UCC), Vic, Spain; ² Institute for Research and Innovation in Life Sciences and Health in Central Catalonia (IRIS-CC), Vic, Spain; ³ Department of Health Promotion and Rehabilitation, Institute of Sport Science and Innovations, Lithuanian Sports University, Kaunas, Lithuania; ⁴ Department of Collective Health, Federal University of Rio Grande do Norte, Natal, Brazil; ⁵ Research Centre for Health (ReaCH), School of Health and Life Sciences, Glasgow Caledonian University, Glasgow, UK

Abstract

Background: Age, multimorbidity, immunodeficiency and frailty of older people living in nursing homes make them vulnerable to COVID-19 and overall mortality.

Objective: To estimate overall and COVID-19 mortality parameters and analyse their predictive factors in older people living in nursing homes over a 2-year period.

Method: *Design:* A 2-year prospective longitudinal multicentre study was conducted between 2020 and 2022.

Setting: This study involved five nursing homes in Central Catalonia (Spain).

Participants: Residents aged 65 years or older who lived in the nursing homes on a permanent basis.

Measurements: Date and causes of deaths were recorded. In addition, sociodemographic and health data were collected. For the effect on mortality, survival curves were performed using the Kaplan-Meier method and multivariate analysis using Cox regression.

Results: The total sample of 125 subjects had a mean age of 85.10 years (standard deviation = 7.3 years). There were 59 (47.2%) deaths at 24 months (95% confidence interval, CI, 38.6–55.9) and 25 (20.0%) were due to COVID-19, mostly in the first 3 months. In multivariate analysis, functional impairment (hazard ratio, HR 2.40; 95% CI 1.33–4.32) was a significant risk factor for mortality independent of age (HR 1.17; 95% CI 0.69–2.00) and risk of sarcopenia (HR 1.40; 95% CI 0.63–3.12).

Conclusion: Almost half of this sample of nursing home residents died in the 2-year period, and one fifth were attributed to COVID-19. Functional impairment was a risk factor for overall mortality and COVID-19 mortality, independent of age and risk of sarcopenia.

Keywords

Predictive factors · Survival · Aged · Long-term care · Pandemic



Scan QR code & read article online




Fig. 1 ▲ Flow chart of the sampling process

Abbreviations

<i>AHT</i>	Arterial hypertension
<i>BMI</i>	Body mass index
<i>CFS</i>	Clinical frailty scale
<i>CI</i>	Confidence interval
<i>COVID-19</i>	Coronavirus disease
<i>CVA</i>	Cerebral vascular accident
<i>HR</i>	Hazard ratio
<i>ICC</i>	Interclass correlation coefficient
<i>MDS</i>	Minimum data set
<i>MNA</i>	Mini nutritional assessment
<i>NH</i>	Nursing homes
<i>PCR</i>	C-reactive protein
<i>PPE</i>	Personal protective equipment
<i>RAT</i>	Rapid antigen tests
<i>SARC-F</i>	Questionnaire assistance in walking, rising from chair, climbing stairs and falling
<i>SB</i>	Sedentary behaviour
<i>SD</i>	Standard deviation
<i>SPSS</i>	Statistical Package Social Sciences
<i>STROBE</i>	Strengthening the Reporting of Observational Studies in Epidemiology
<i>UI</i>	Urinary incontinence
<i>WTMB</i>	Waking-time movement behaviours

Introduction

The coronavirus disease 2019 (COVID-19) pandemic began to have a major impact on society in 2019 [1] having unprecedented consequences on global health and economic systems.

In developed European countries with a very high older population the COVID-19 mortality was 83.7% for people > 70 years and 16.2% for people younger than 69 years in 2020 [2] with a higher prevalence of COVID-19 deaths in nursing homes (NH) [3]. Health problems and geriatric syndromes associated with ageing also determined the risk of mortality [4–9].

At the beginning of the pandemic Spain had a total of 326,613 people institutionalised in NHs [10] and a study conducted in Madrid reported a 14% mortality rate in older adults with COVID-19 in NHs [7]. The COVID-19 mortality has already been extensively studied in several countries, although most of these follow-ups have

not exceeded 1 year [11]. Longer follow-ups would enable more accurate data and the identification of predictive factors that may be relevant to clinical practice.

The main aim of the study was to estimate overall and COVID-19 mortality parameters and analyse their predictive factors in older people living in NHs over a 2-year period.

Methodology

Study design and population

This is a 2-year multicentre observational cohort study. The study was conducted in five NH in Central Catalonia, Spain. It was designed following the Strengthening the Reporting of Observational studies in Epidemiology (STROBE) standards for cohort studies [12, 13]. Residents aged 65 years and older permanently living in NH were included. Those in a coma or palliative care (short-term prognosis), those who refused (or their legal guardian) to participate in the study and those who left the NH during the 2-year cohort period were excluded.

Sample size

For the calculation of the sample, the article by Burgaña Agoües et al. (2021) was taken as a reference due to its methodological similarity to the present article: Burgaña studied pandemic mortality due to COVID-19 in Spain, in people over 65 years of age resident in NH. Considering the findings of Burgaña Agoües et al. (2021) [14], i.e. the difference in proportions between individuals with severe functional impairment (23.0%) and deceased (11.1%), and with a confidence interval (CI) of 95% and a power of 80%, a sample size of 122 participants was estimated.

Study procedures

The primary outcomes were all-cause mortality and COVID-19 over the 2-year follow-up period. The mortality registry included cause and date of death, deaths in total, deaths due to COVID-19 including confirmed cases, and deaths due to symptomatic suspicion of COVID-19 [15]. Additional COVID-19 data collected included the presence of the disease,

Table 1 Descriptive analysis of the sample of institutionalized older adults living in nursing homes of Central Catalonia, Spain (<i>n</i> = 125)	
Variables	Frequency (%) / mean (SD)
<i>NH type</i>	
State subsidized places	40 (32.0)
Private	85 (68.0)
Smoking	6 (4.8)
Drinkers	9 (12.7)
<i>Chronic conditions</i>	
Hypertension	80 (64.0)
Dementia	68 (54.4)
Cardiac pathology	51 (40.8)
Depression	36 (28.8)
Diabetes mellitus type 2	36 (28.8)
Renal pathology	32 (25.6)
CVA	25 (20.0)
Cancer	23 (18.4)
Pulmonary pathology	22 (17.6)
<i>Urinary incontinence</i>	
Yes	87 (69.6)
No	35 (28.0)
Unclassifiable	3 (2.4)
<i>Fecal incontinence</i>	
Yes	36 (28.8)
No	87 (69.6)
Unclassifiable	2 (1.6)
Fall/s in previous year	58 (46.4)
<i>Nutritional status</i>	
Involuntary weight loss	25 (13.5)
Risk of malnutrition or malnourished	56 (44.8)
Obesity	78 (62.4)
Risk of sarcopenia	94 (75.2)
<i>Functional impairment</i>	
Independent	7 (5.6)
Slightly dependent	47 (37.6)
Moderately dependent	19 (15.2)
Severely or totally dependent	52 (41.6)
<i>Cognition</i>	
No cognitive deficit	10 (8.0)
Mild cognitive deficit	20 (16.0)
Moderate cognitive deficit	24 (19.2)
Severe cognitive deficit	58 (46.4)
Unknown	13 (10.4)
<i>SB and WTMB</i>	
Waking hours	11.0 (1.5 SD)
% of waking time in SB	82.6% (17.5 SD)
Hours in upright position (standing and walking)	1.6 (1.7 SD)
Steps per day	1.345 (2417.4 SD)
Sit to stand transitions per day	18.2 (18.3 SD)
Hospitalization	26 (0.43 SD)
<i>SD</i> standard deviation, <i>NH</i> nursing homes, <i>CVA</i> cerebral vascular accident, <i>SB</i> sedentary behaviour, <i>WTMB</i> waking-time movement behaviour	

whether they had symptoms [16], the performance of COVID-19 screening tests such as C-reactive protein (CRP), serological tests, and/or rapid antigen tests for SARS-CoV-2 (RAT) [17]. The information was collected through on-line interviews with NH professionals as due to COVID-19, they could not be accessed.

Sociodemographic and health information was obtained from health centre records and cross-checked with health professionals. All sociodemographic variables and those described in this article were collected at baseline (January 2020), just before the onset of the COVID-19 pandemic in Spain. Falls (number) during the last year were obtained from NH registers. Nutritional status was assessed using the mini nutritional assessment (MNA) [18]. The SARC-F [19] was used to identify individuals at risk of developing sarcopenia. Functional capacity was measured using the modified Barthel index and results were classified according to the degree of dependency as: independent, slightly dependent, moderately dependent, severely or totally dependent [20]. Continence status was reported using section H of the minimum data set (MDS) version 3.024 [21]. Cognitive status was assessed using the Pfeiffer scale [22] and frailty using the clinical frailty scale (CFS) [23]. Sedentary behaviour (SB) and waking-time movement behaviours (WTMB) were assessed using the ActivPAL 3TM activity monitor (PAL Technologies Ltd., Glasgow, UK) [24].

The study was conducted over a 2-year period and ended in March 2022.

Statistical analysis

The nominal and ordinal quantitative variables were expressed according to frequency in percentages and the quantitative variables with mean and standard deviation (SD). Survival curves were formed using the Kaplan-Meier method and multivariate analysis was performed by Cox regression, using the hazard ratio (HR) as the measure of effect. The Statistical Package for the Social Sciences 27 (SPSS Inc., Chicago, IL, USA) was used for the analysis.

Table 2 Association of variables to health, functional and sociodemographic ¹ with mortality (n = 125)					
Variables	n (%)	Number of deaths	Number of survivors	Probability of death (%)	p (Log rank)
<i>Functional impairment (n = 125)</i>					
No/mild or moderate	73 (58.4)	24	49	65.3	0.001*
Total impairment	52 (41.6)	35	17	31.6	
<i>Urinary incontinence (n = 122)</i>					
No	35 (28.0)	10	25	66.8	0.008*
Yes	87 (69.6)	49	38	43.2	
<i>Risk of sarcopenia (n = 125)</i>					
No	31 (24.8)	9	22	71.0	0.018*
Yes	94 (75.2)	50	44	45.0	
<i>% waking time in SB (n = 84)</i>					
≤ 85%	41 (48.8)	11	30	73.2	0.028*
> 85%	43 (51.2)	22	21	48.4	
<i>Fecal incontinence (n = 123)</i>					
No	87 (70.7)	35	52	57.9	0.029*
Yes	36 (29.3)	22	14	38.4	
<i>NH type (n = 125)</i>					
State subsidized places	40 (32.0)	14	26	64.8	0.076
Private	85 (68.0)	45	40	46.1	
<i>Nutritional status (n = 79)</i>					
Normal	23 (29.2)	7	16	69.6	0.082
At risk or malnourished	56 (70.8)	31	25	43.6	
<i>Drinkers (n = 71)</i>					
No	62 (87.3)	34	32	45.5	0.249
Yes	9 (12.7)	1	4	80.0	
SB sedentary behaviour, NH nursing homes					
¹ With a p value lower than 0.250					
* Statistically significant (< 0.05)					

Results

We recruited 125 people, 67.6% of the total number of NH residents in the main study. Finally, 7 (3.8%) participants who left NHs to reside elsewhere were excluded (▣ Fig. 1).

The mean age of the participants was 85.1 years (SD = 7.3 years) and 104 (83.2%) were female. The mean number of months living in NH was 27.5 months (SD = 112.14 months). The analysis of health and sociodemographic variables is described in ▣ Table 1.

In the 2-year period from baseline to the end of the study, 59 participants (47.2%) died, of whom 25 (20.0%) died from COVID-19 and 34 (27.2%) from other causes. All COVID-19 deaths occurred in the first year of the study: 44 (74.5%) of the 59 individuals had already died within the first 90 days (at the peak of COVID-19).

Survival and associated factors according to the variable mortality

In the bivariate analysis, mortality was associated with functional impairment, urinary incontinence (UI), faecal continence, risk of sarcopenia, % of waking time in SB and with a p-value of less than 0.05. All other health and sociodemographic variables were not significant (▣ Table 2).

The variables were tested for collinearity and none of them showed collinearity with each other. A multivariate analysis was performed with the model with adjusted values including the variables age with severe functional impairment and risk of sarcopenia. The result showed that functional impairment predicted mortality independently of age (which was not statistically significant) and sarcopenia risk (▣ Table 3).

Survival and associated factors according to the variable COVID-19 or other-cause mortality

In the univariate analysis, functional impairment, living in a private NH, being older than 86 years, malnutrition and being female were risk factors for COVID-19 mortality. Functional impairment was associated with mortality from other health causes with a p-value of less than 0.050 (▣ Tables 4 and 5; ▣ Fig. 2).

We tested for collinearity between the variables and none of them were collinear with each other. The number of individuals in this variable is 59. Different combinations of significant variables, such as functionality, age and type of NH among others, were tested by multivariate analysis, but no significant results were found (▣ Table 5).

Table 3 Survival univariate and multivariate Cox analysis ¹ in older people living in NHs (n = 125)						
Variables	HR (ref.)	CI (95%)	p (Cox)	HR (ref.)	CI (95%)	p (Cox)
	Univariate analysis			Multivariate analysis		
<i>Functional impairment (n = 125)</i>						
No/mild or moderate	–	–	–	–	–	–
Total impairment	2.77	(1.64–4.67)	0.001*	2.40	(1.33–4.32)	0.003*
<i>Urinary incontinence (n = 122)</i>						
Yes	–	–	–	–	–	–
No	2.43	(1.23–4.81)	0.010*	–	–	–
<i>Risk of sarcopenia (n = 125)</i>						
Yes	–	–	–	–	–	–
No	2.29	(1.12–4.66)	0.022*	1.40	(0.63–3.12)	0.403
<i>Fecal incontinence (n = 123)</i>						
No	–	–	–	–	–	–
Yes	1.80	(1.05–3.07)	0.031*	–	–	–
<i>% waking time in SB (n = 84)</i>						
≤ 85%	–	–	–	–	–	–
> 85%	2.20	(1.07–4.55)	0.033*	–	–	–
<i>NH type (n = 125)</i>						
Private	–	–	–	–	–	–
State subsidized places	0.85	(0.32–1.06)	0.080	–	–	–
<i>Nutritional status (n = 79)</i>						
Normal	–	–	–	–	–	–
Risk or malnourished	2.04	(0.89–4.64)	0.089	–	–	–
<i>Age (years, n = 125)</i>						
≤ 85	–	–	–	–	–	–
> 86	1.33	(0.78–2.26)	0.284	1.17	(0.69–2.00)	0.549

HR hazard ratio, CI confidence interval, SB sedentary behaviour, NH nursing homes
¹ Variables with a p value lower than 0.250 in univariate analysis are shown
* Statistically significant (p < 0.05)




Fig. 2 ◀ Kaplan-Meier survival estimates

Discussion

The main objective of this study was to examine the incidence of all-cause and COVID-19 mortality and to analyse the predictive factors in older NH residents

over a 2-year period since the onset of the pandemic.

The results indicate that almost half of participants died with 20% being attributed to COVID-19. Most of the deaths (74.5%) were in the first 3 months of the study, coinciding with the outbreak of the

COVID-19 pandemic in Spain. In the second year of the study, the survival curve became horizontal again, after the implementation of preventive measures. This study shows a higher incidence of mortality in the 1-year period than other studies. Several articles on the pandemic phase report data on excess mortality [11]. A study in Barcelona reported a 3-month COVID-19 mortality rate of 11.1% in institutionalised older people [14]. For deaths from other causes, they reported excess mortality among institutionalised cases (34.8%) [14].

We also report the association of health, social and demographic variables with mortality: functional impairment, UI, sarcopenia risk and % of waking time in SB were found to be factors associated with mortality and functional impairment, type of NH and age were associated with COVID-19 mortality. The literature shows that UI and risk of sarcopenia are asso-

Table 4 Analysis of the association of health and sociodemographic variables¹ NH residents with mortality (COVID-19 or other causes) (n = 59)

Variables	n (%)	Deaths for covid-19 (number of events)	Deaths due to other causes (number of events)	Probability of death (%)	p (Log rank)
<i>Nutritional status (n = 38)</i>					
Normal	7 (18.4)	5	2	28.6	0.022*
Risk or malnourished	31 (81.6)	10	21	61.8	
<i>Sex (n = 59)</i>					
Male	9 (15.2)	6	3	14.6	0.041*
Female	50 (84.7)	19	31	29.8	
<i>Functional impairment (n = 59)</i>					
No/mild or moderate	24 (40.6)	15	9	26.1	0.045*
Total impairment	35 (59.3)	10	25	63.5	
<i>% waking time in SB (n = 33)</i>					
≤ 85%	11 (33.3)	7	4	36.4	0.082
> 85%	22 (66.6)	6	16	68.7	
<i>Fecal incontinence (n = 57)</i>					
No	35 (61.4)	18	17	37.6	0.132
Yes	22 (38.6)	6	16	68.2	
<i>Hospitalizations (n = 46)</i>					
No	34 (73.9)	17	17	32.0	0.194
Yes	12 (26.1)	4	8	58.2	

SB sedentary behaviour, NH nursing homes
¹ With a p value lower than 0.250
 * Statistically significant (< 0.05)

Table 5 Univariate Cox analysis: association of COVID-19 mortality with other causes of death in relation to covariables¹ in older people living in NHs (n = 59)

Variables	HR ref. (exp ^B)	CI (95%)	p value
<i>(Univariate analysis)</i>			
<i>Functional impairment (n = 59)</i>			
No/mild or moderate impairment	–	–	–
Total impairment	2.04	(1.40–2.97)	0.001*
<i>NH type (n = 59)</i>			
State subsidized places	–	–	–
Private	1.73	(1.13–2.66)	0.012*
<i>Age (years, n = 59)</i>			
≤ 85	–	–	–
≥ 86	1.47	(1.03–2.11)	0.035*
<i>Fecal incontinence (n = 57)</i>			
No	–	–	–
Yes	1.47	(0.99–2.18)	0.055
<i>Urinary incontinence (n = 59)</i>			
Yes	–	–	–
No	1.41	(0.95–2.08)	0.090
<i>% waking time in SB (n = 33)</i>			
≤ 85%	–	–	–
> 85%	1.36	(0.88–2.09)	0.161

HR hazard ratio, CI confidence interval, NH nursing homes, SB sedentary behaviour
¹ Variables with a p value lower than 0.250 are shown
 * Statistically significant (< 0.05)

ciated with mortality [25–27] and those who spent more time in SB had a higher risk of mortality [28].

Unlike other studies, our data show an increase of mortality in private NHs [29]. The NH type and size influenced mortality during the COVID-19 pandemic in other studies [29]. Braun et al. (2020) attributed these results to the lack of organisation and shortage of personal protective equipment (PPE) in private NHs [29].

This study has the limitation of the COVID-19 pandemic, which impeded access to NHs, increased deaths in older people and did not enable us to have a larger sample.

The strength of the study lies in the telematic data collection in NHs. This enabled us to extract real information on the health and social status of residents. The fact that we collected data prior to the start of the pandemic allowed us to make a comparison of the health and sociodemographic status of institutionalised older people. By having a cross-sectional analysis of the prepandemic sample, we provide data on the factors that predicted the risk of dying in a 2-year follow-up.

Conclusion

Almost half of this sample of NH residents died during the 2-year observation period. One fifth of deaths were attributed to COVID-19 mostly in the first quarter, coinciding with the peak of the pandemic. Functional impairment was a risk factor for overall mortality and COVID-19 mortality, independent of age and risk of sarcopenia.

Corresponding address

Javier Jerez-Roig

Research group on Methodology, Methods, Models and Outcomes of Health and Social Sciences (M3O). Faculty of Health Sciences and Welfare. Centre for Health and Social Care Research (CESS), University of Vic-Central University of Catalonia (UVic-UCC) C. Sagrada Família 7, 08500 Vic, Spain
Javier.jerez@uvic.cat

Acknowledgements. We thank all the members of the study NH for their contribution to this work and the older adults who participated in the study.

Funding. This work was supported by the Hestia Chair from Universitat Internacional de Catalunya (grant number BI-CHAISS-2019/003) and the research grant from the Catalan Board of Physiotherapists Code R03/19.

Dyego Leandro Bezerra de Souza thank CNPq (Brazilian National Council for Scientific and Technological Development) productivity grants 315962/2023-2.

Author Contribution. Anna Escribà-Salvans: conceptualization, methodology, formal analysis, investigation, writing—original draft, writing—review and editing. Eduard Minobes-Molina: conceptualization, methodology, writing—review and editing, funding acquisition. Pau Farrés-Godayol: formal analysis, writing—review and editing. Dyego Leandro Bezerra de Souza: writing—review and editing. Dawn A Skelton: supervision, writing—review and editing. Javier Jerez-Roig: conceptualization, methodology, supervision, funding acquisition.

Funding. Open Access funding provided thanks to the CRUE-CSIC agreement with Springer Nature.

Availability of data and material. The datasets used and/or analyzed during the current study are available from the corresponding author on reasonable request.

Declarations

Conflict of interest. A. Escribà-Salvans, J. Jerez-Roig, P. Farrés-Godayol, D.L. Bezerra de Souza, D.A. Skelton and E. Minobes-Molina declare that they have no competing interests.

Ethical approval was obtained from the Ethics and Research Committee of the University of Vic—Central University of Catalonia (registration number 92/2019 and 109/2020). Signed informed consent was obtained from the residents or the legal guardians. The project meets the criteria required in the Helsinki Declaration as well as the Organic Law 3/2018 (5 December) on the Protection of Personal Data and Guarantee of Digital Rights.

Open Access. This article is licensed under a Creative Commons Attribution 4.0 International License, which permits use, sharing, adaptation, distribution and reproduction in any medium or format, as long as you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons licence, and indicate if changes were made. The images or other third party material in this article are included in the article's Creative Commons licence, unless indicated otherwise in a credit line to the material. If material is not included in the article's Creative Commons licence and your intended use is not permitted by statutory regulation or exceeds the permitted use, you will need to obtain permission directly from the copyright holder. To view a copy of this licence, visit <http://creativecommons.org/licenses/by/4.0/>.

References

- Licher S, Terzikhan N, Splinter MJ et al (2021) Design, implementation and initial findings of COVID-19 research in the Rotterdam Study: leveraging existing infrastructure for population-based investigations on an emerging disease. *Eur J Epidemiol* 36(6):649–654
- Abbatecola AM, Antonelli-Incalzi R (2020) Editorial: Espiral de fragilidad COVID-19 en pacientes italianos mayores. *Rev Nutr Salud Envejec* 24(5):453–455. <https://doi.org/10.1007/s12603-020-1357-9>
- Levin AT, Owusu-Boaitey N, Pugh S et al (2022) Assessing the burden of COVID-19 in developing countries: systematic review, meta-analysis and public policy implications. *BMJ Glob Health* 7(5):e8477. <https://doi.org/10.1136/bmjgh-2022-008477>
- Beard JR, Officer A, de Carvalho IA et al (2016) The World report on ageing and health: a policy framework for healthy ageing. *Lancet* 387(10033):2145–2154. [https://doi.org/10.1016/S0140-6736\(15\)00516-4](https://doi.org/10.1016/S0140-6736(15)00516-4)
- Damián J, Pastor-Barriuso R, García López FJ et al (2017) Urinary incontinence and mortality among older adults residing in care homes. *J Adv Nurs* 73(3):688–699. <https://doi.org/10.1111/jan.13170>
- Escribà-Salvans A, Rierola-Fochs S, Farrés-Godayol P et al (2023) Risk factors for developing symptomatic COVID-19 in older residents of nursing homes: A hypothesis-generating observational study. *JFSF* 8(2):74–82. <https://doi.org/10.22540/JFSF-08-074>
- Moreno-Torres V, de la Fuente S, Mills P, Muñoz A, Muñoz E, Ramos A, Fernández-Cruz A, Arias A, Pintos I, Vargas JA, Cuervas-Mons V, de Mendoza C (2021) Major determinants of death in patients hospitalized with COVID-19 during the first epidemic wave in Madrid, Spain. *Ann Univ Mariae Curie Skłodowska [Med]* 100(16):E25634. <https://doi.org/10.1097/MD.00000000000025634>
- Regato-Pajares P, Villacañas-Novillo E, López-Higuera MJ et al (2023) Primary care and elderly people in nursing homes: proposals for improvement after the experience during the pandemic. *Rev Clin Med Fam* 16(1):24–32. <https://doi.org/10.55783/rcmf.160105>
- Fallon A, Dukelow T, Kennelly SP, O'Neill D (2020) COVID-19 in nursing homes. *QJM* 113(6):391–392. <https://doi.org/10.1093/qjmed/hcaa136>
- Rodríguez Rodríguez P, Gonzalo Jiménez E (2022) COVID-19 in nursing homes: structural factors and experiences that support a change of model in Spain. *Gaceta Sanit* 36(3):270–273. <https://doi.org/10.1016/j.gaceta.2021.09.005>
- Cases L, Vela E, Santaugènia González SJ et al (2023) Excess mortality among older adults institutionalized in long-term care facilities during the COVID-19 pandemic: a population-based analysis in Catalonia. *Front Public Health* 11:1208184. <https://doi.org/10.3389/fpubh.2023.1208184>
- von Elm E, Altman DG, Egger M et al (2008) The strengthening of reporting of observational studies in epidemiology (STROBE) statement: guidelines for reporting observational studies. *J Clin Epidemiol* 61(4):344–349. <https://doi.org/10.1016/j.jclinepi.2007.11.008>
- Farrés-Godayol P, Jerez-Roig J, Minobes-Molina E et al (2021) Urinary incontinence and sedentary behaviour in nursing home residents in Osona, Catalonia: protocol for the OsoNaH project, a multicentre observational study. *BMJ Open* 11(4):e41152. <https://doi.org/10.1136/bmjopen-2020-041152>
- Burgaña Agouès A, Serra Gallego M, Hernández Resa Retal (2021) Risk factors for COVID-19 morbidity and mortality in Institutionalised elderly people. *Int J Environ Res Public Health* 18(19):10221. <https://doi.org/10.3390/ijerph181910221>
- Comas-Herrera A, Zalakain J Mortality associated with COVID-19 outbreaks in care homes: early international evidence. Resources to support community and institutional Long-Term Care responses to COVID-19. 2020;(April):1–6. <https://ltccovid.org/2020/04/12/mortality-associated-with-covid-19-outbreaks-in-care-homes-early-international-evidence/>
- Dhama K, Patel SK, Natesan S et al (2020) COVID-19 in the elderly people and advances in vaccination approaches. *Hum Vaccin Immunother* 16(12):2938–2943. <https://doi.org/10.1080/21645515.2020.1842683>
- López de la Iglesia J, Fernández-Villa T, Rivero A et al (2020) Predictive factors of COVID-19 in patients with negative RT-qPCR. *Sermergen* 46(Suppl 1):6–11. <https://doi.org/10.1016/j.semserg.2020.06.010>
- Dewar AM, Thornhill WA, Read LA (1994) The effects of tefluthrin on beneficial insects in sugar beet. In: The mini nutritional assessment (MNA) and its use in grading the nutritional state of elderly patients, pp 987–992
- Malmstrom TK, Miller DK, Simonsick EM et al (2016) SARC-F: a symptom score to predict persons with sarcopenia at risk for poor functional outcomes. *J Cachexia Sarcopenia Muscle* 7(1):28–36. <https://doi.org/10.1002/jcsm.12048>
- Cid-Ruzafa J, Damián-Moreno J (1997) Disability evaluation: Barthel's index. *Rev Esp Salud Publica* 71(2):127–137 (Erratum in: *Rev Esp Salud Publica* 1997 Jul-Aug;71(4):411)
- Klusch L (2012) SMD 3.0 and its impact on bladder and bowel care. *Provider* 38(6):33–37
- Martínez de la Iglesia J, Dueñas Herrero R, Onís Vilches MC et al (2001) Spanish language adaptation and validation of the Pfeiffer's questionnaire (SPMSQ) to detect cognitive deterioration in

- people over 65 years of age. *Med Clin (Barc)* 117(4):129–134. [https://doi.org/10.1016/s0025-7753\(01\)72040-4](https://doi.org/10.1016/s0025-7753(01)72040-4)
23. Rockwood K, Song X, MacKnight C et al (2005) A global clinical measure of fitness and frailty in elderly people. *CMAJ* 173(5):489–495. <https://doi.org/10.1503/cmaj.050051>
 24. Grant PM, Dall PM, Mitchell SL et al (2008) Activity-monitor accuracy in measuring step number and cadence in community-dwelling older adults. *J Aging Phys Act* 16(2):201–214. <https://doi.org/10.1123/japa.16.2.201>
 25. Huang P, Luo K, Wang C et al (2021) Urinary incontinence is associated with increased all-cause mortality in older nursing home residents: a meta-analysis. *J Nurs Scholarsh* 53(5):561–567. <https://doi.org/10.1111/jnu.12671>
 26. Yoshioka T, Kamitani T, Omae K et al (2021) Urgency urinary incontinence, loss of independence, and increased mortality in older adults: A cohort study. *PLoS ONE* 16:1–13. <https://doi.org/10.1371/journal.pone.0245724>
 27. Yang M, Jiang J, Zeng Y et al (2019) Sarcopenia for predicting mortality among elderly nursing home residents. *Medicine* 98(7):e14546. <https://doi.org/10.1097/MD.00000000000014546>
 28. Ku PW, Steptoe A, Liao Y et al (2019) A threshold of objectively-assessed daily sedentary time for all-cause mortality in older adults: a meta-regression of prospective cohort studies. *J Clin Med* 8(4):564. <https://doi.org/10.3390/jcm8040564>
 29. Braun RT, Yun H, Casalino LP et al (2020) Comparative performance of private equity-owned US nursing homes during the COVID-19 pandemic. *JAMA Netw Open* 3(10):e2026702. <https://doi.org/10.1001/jamanetworkopen.2020.26702>

Publisher's Note. Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.

Gesundheitliche und soziodemografische Determinanten der Übersterblichkeit in spanischen Pflegeheimen während der COVID-19-Pandemie: eine 2-jährige prospektive Längsschnittstudie

Hintergrund: Alter, Multimorbidität, Immunschwäche und Gebrechlichkeit älterer Menschen, die in Pflegeheimen leben, machen sie anfällig für COVID-19 und erhöhen die Gesamtmortalität.

Zielsetzung: Schätzung der Gesamt- und COVID-19-Mortalität und Analyse ihrer prädiktiven Faktoren bei älteren Menschen, die in Pflegeheimen leben, über einen Zeitraum von 2 Jahren.

Methode: Diese prospektive, multizentrische Längsschnittstudie wurde zwischen 2020 und 2022 in 5 Pflegeheimen in Zentral-Katalonien (Spanien) durchgeführt. Teilnehmer waren Bewohner im Alter von 65 Jahren oder älter, die dauerhaft in den Pflegeheimen lebten. Datum und Ursachen der Todesfälle wurden erfasst. Darüber hinaus wurden soziodemografische und gesundheitliche Daten erhoben. Für die Auswirkung der Sterblichkeit wurden Überlebenskurven nach der Kaplan-Meier-Methode und eine multivariate Analyse mittels Cox-Regression durchgeführt.

Ergebnisse: Die Gesamtstichprobe von 125 Personen hatte ein Durchschnittsalter von 85,10 Jahren (Standardabweichung = 7,3). Nach 24 Monaten traten 59 (47,2 %) Todesfälle auf (95 % Konfidenzintervall [KI] 38,6–55,9), von denen 25 (20,0 %) auf COVID-19 zurückzuführen waren, zumeist in den ersten 3 Monaten. In der multivariaten Analyse war die Funktionseinschränkung (Hazard-Ratio [HR]: 2,40; 95 % KI 1,33–4,32) ein signifikanter Risikofaktor für die Sterblichkeit, unabhängig vom Alter (HR: 1,17; 95 % KI 0,69–2,00) und dem Risiko der Sarkopenie (HR: 1,40; 95 % KI 0,63–3,12).

Schlussfolgerung: Fast die Hälfte dieser Stichprobe von Pflegeheimbewohnern starb innerhalb von 2 Jahren, ein Fünftel davon wurde auf COVID-19 zurückgeführt. Funktionseinschränkungen waren ein Risikofaktor für die Gesamt- und COVID-19-Mortalität, unabhängig von Alter und Sarkopenierisiko.

Schlüsselwörter

Vorhersagefaktoren · Überleben · Alter · Langzeitpflege · Pandemie